

Wengen - Alpine Flowers of the Swiss Alps

Naturetrek Tour Report

24 June - 1 July 2018



Cypripedium calceolus



Jungfrau



Rhododendron hirsutum



Viola cenisia

Report and images by David Tattersfield



Naturetrek Mingledown Barn Wolf's Lane Chawton Alton Hampshire GU34 3HJ UK

T: +44 (0)1962 733051

E: info@naturetrek.co.uk

W: www.naturetrek.co.uk

Tour participants: David Tattersfield (leader) with 15 Naturetrek clients.

Day 1

Sunday 24th June

After assembling at Zurich airport, we were soon enjoying the comfort of the inter-city train to Bern. We changed trains here and again at Interlaken, where we joined the regional train to Lauterbrunnen. The last leg of the journey was on the delightful cogwheel railway to Wengen, where we arrived around 8.00pm. It was just a short walk to our hotel, where a delicious evening meal was waiting for us. After talking through the plans for the exciting week ahead, we retired to bed.

Day 2

Monday 25th June

We woke to a cloudy and cool day, with high cloud over the summits. We took the Mannlichen cable-car, high above the village, and set off to explore a wide range of habitats, as we made our way, slowly, towards Mannlichen summit, at 2343 metres. We found a rich flora, on shaded banks and cliffs, typical of areas where snow lies late into the season. A range of dwarf shrubs included Net-leaved Willow, *Salix reticulata*, Retuse-leaved Willow *Salix retusa* and Mountain Avens *Dryas octopetala*, with splashes of colour provided by Bird's-eye Primrose *Primula farinosa*, Moss Campion *Silene acaulis*, Alpine Cinquefoil *Potentilla crantzii* and Spring Gentian *Gentiana verna*. As we climbed higher, we found a colourful tapestry of flowers in the turf, from the purple Whorled Lousewort *Pedicularis verticillata* and blue Alpine Forget-me-not *Myosotis alpestris* to the rich yellows of Globeflower *Trollius europaeus* and Alpine Pasque-flower *Pulsatilla alpina* subsp. *apiifolia*. Alpine Aster *Aster alpinus* and deep-purple Alpine Sainfoin *Hedysarum hedysaroides* framed the view of the deep, glacial trench of the Lauterbrunnen Valley, far below us. Near the edges of the last remaining snow, there were still one of two white flowers of Spring Crocus *Crocus vernus* and Kuepfer's Buttercup *Ranunculus kuepferi* and the fringed lavender bells of Alpine Snowbell *Soldanella alpina*. Just beyond the summit, we found Crimson-tipped Lousewort *Pedicularis oederi*, growing on the north-facing slopes.

We regrouped at the Mannlichen Berghaus, before having lunch on a grassy knoll overlooking Grindelwald. For the rest of the afternoon, we followed an easy path down to Kleine Scheidegg, with the brooding, north faces of the Eiger, Monch and Jungfrau appearing, occasionally, through gaps in the cloud. There were many more plants to find and identify and we made slow progress, eventually catching the 5.30pm train back to Wengen.

Day 3

Tuesday 26th June

On a warm on sunny morning, we took the short train ride to Wengernalp, where we started by exploring the vicinity of the station platform. After a little search, we located a few Frog Orchids *Coeloglossum viride*, their muted colours blending well with their background. Over coffee, we were able to pick out a herd of Alpine Ibex, grazing below the cliffs on the opposite side of the valley. We walked down through colourful meadows to Wixi, where we found the deep-blue Bavarian Gentian *Gentiana bavarica*, growing in wet flushes, together with lots of Broad-leaved Marsh-orchid *Dactylorhiza majalis*, with flowers and bracts of an intense deep purple. As we descended to Biglenalp, some of the special plants we encountered included a single specimen of *Gymnadenia x suaveolens* (a hybrid between Fragrant Orchid and Vanilla Orchid), Serrated Wintergreen *Orthilia secunda*, One-flowered Wintergreen *Moneses uniflora* and the parasitic Coralroot Orchid *Corallorhiza trifida*.

We chose a sunny bank, close to the stream, to have our picnic, an idyllic spot, with cows grazing contentedly, and views of the glistening ice-fields of the Jungfrau. Nearby, were the beautiful large blue flowers of *Aquilegia alpina* and a single plant of Lady's-slipper *Cypripedium calceolus*. Our return route took us past the farmstead of Mettlenalp, on a lovely walk through forest and pastures, leading, eventually, back to Wengen. Most of the group took advantage of a train from Almend, and were back at around 5.30pm. Some of us took a longer route and explored the forest above Wengen, where we found a population of St Bernard's Lily *Anthericum liliago*. Over dinner, we enjoyed a lovely alpenglow over the Jungfrau.

Day 4

Wednesday 27th June

We walked down to Lauterbrunnen, finding a new range of plants to interest us. In the forest were Herb-Paris *Paris quadrifolia* and Whorled Solomon's-seal *Polygonatum verticillatum* and there were beautiful patches of Dark Columbine *Aquilegia atrata* and the large Cabbage Thistle *Cirsium oleraceum*, in the meadows. We arrived in Lauterbrunnen at 11.50am and had plenty of time for a coffee break, before taking the bus to Trummelbach. We had lunch at a convenient picnic table, by the river, before visiting the Trummelbach Falls. The sights and sounds of the waterfalls were a memorable experience, and we marvelled at the ingenuity of the Swiss engineers who had constructed the access tunnels, one hundred years previously. New plants, around the falls included Branched St Bernard's Lily *Anthericum ramosum*, Blue Saxifrage *Saxifraga caesia*, and the rhizomatous Mountain Bladder-fern *Cystopteris montana*. Most of the group walked back to Lauterbrunnen, along the valley and caught a train back to Wengen.

Ivan and I returned by a more energetic route, making the 900metre climb to the top of the cliffs, above the falls, past the little farmstead at Brech, before descending, on the previous day's track, to Wengen. It was an exhilarating walk with some incredible views and botanical highlights which included Lesser Butterfly-orchid *Platanthera bifolia*, Large Yellow Foxglove *Digitalis grandiflora* and Red Helleborine *Cephalanthera rubra*. We were back at the hotel within four hours, well satisfied with our efforts.

Day 5

Thursday 28th June

We had another sunny day, with cloud building over the mountains in the afternoon, a pattern that was repeated on every day. We took the train down to Lauterbrunnen and the cable-car to Grutschalp, where we boarded another train to Murren, high on the edge of the cliff, opposite the Trummelbach Falls. After a short stroll through the village, we took the funicular railway to the restaurant at Almendhubel and had a coffee break. The vegetation on this side of the valley provided a complete contrast to what we had seen previously. Tall herbs predominated, among thickets of Green Alder *Alnus viridis* and a luxuriance of ferns. Mountain Pansy *Viola lutea* was common, in a variety of colours, and marshy ground was thick with Broad-leaved Marsh-orchid *Dactylorhiza majalis*.

After lunch, we followed the Mountain View Trail, across heather-clad slopes and lush pasture, with an abundance of Aconite-leaved Buttercup *Ranunculus aconitifolius* and the huge pleated leaves of White False-helleborine, *Veratrum album*, both plants toxic to grazing animals. We descended through damp thickets of Green Alder and tall herbs, where we found Alpine Leek *Allium victorialis*, Yellow Pea *Lathyrus laevigatus* subsp. *occidentalis* and large stands of Alpine Blue-sowthistle *Cicerbita alpina*. Our final descent was through the forest, where there were large colonies of Lesser Twaybade, *Neottia cordata* and the curious *Streptopus amplexifolius*, with small,

yellowish, pendant flowers, on bent pedicels, hidden beneath its leaves. Eventually, we reached Grutschalp station and our return cable-car and were back in Wengen for 6.00pm.

Day 6

Friday 29th June

We had a relaxing start to the day, travelling by train down to Wilderswil, and then joining the delightful little train, for the steep climb to Schynige Platte. After coffee, we spent an absorbing morning in the Alpine Garden, enjoying its natural setting and the superb display of plants, each carefully labelled. Many plants were familiar to us, by now, but there were many others that had been brought from other regions of Switzerland, each growing in its preferred habitat. The whole garden is an enormous credit to the dedicated team of gardeners that keep it looking immaculate.

After lunch on the station platform, we climbed to the summit of Gaube and followed the ridge towards the Oberberghorn. The views from up here were tremendous and we looked down on Interlaken and the two lakes, the Brienzensee and Thunersee. We were in a botanical paradise, with superb specimens of Alpine Leek, Yellow Pea and the deep-purple Mountain Milk-vetch *Oxytropis jacquinii*. The limestone slopes of the Oberberghorn were home to a rich diversity of plants, such as the robust, yellow, Leafy Lousewort *Pedicularis foliosa*, Alpine Aster *Aster alpinus* and Wood Pink *Dianthus sylvestris* and Dwarf Buckthorn *Rhamnus pumilus* clung to the rocks above our heads. These were just a few of many that were new to us on the trip. On the limestone screes were extensive colonies of Limestone Fern *Gymnocarpium robertianum*, Angular Solomon's-seal *Polygonatum odoratum*, Whorled Solomon's-seal and Lily-of-the-Valley *Convallaria majalis*. Just enough time remained for us to walk to a second viewpoint, where St Bruno's Lily *Paradisea liliastrum* was common on the grassy slopes, before we had to return to the station, to catch our train. Between trains, at Wilderswil, we enjoyed ice-creams, before the familiar journey back to Wengen. During the evening, over dinner on the terrace, we were treated to our third alpenglow of the week.

Day 7

Saturday 30th June

For our last full day, we took the trains to Kleine Scheidegg and on to the Eigergletscher station and, once again, the weather was perfect. Above the station platform, we made our way past the railway tunnel entrance, onto the moraines, below the west ridge of the Eiger. Although, from a distance they looked bare, closer inspection revealed some wonderful plants among the rocks. Most attractive, among these, were Broad-leaved Mouse-ear *Cerastium latifolium*, Mt Cenis Pansy *Viola cenisia* and the deliciously-scented Round-leaved Penny-cress *Thlaspi rotundifolium*. We followed the narrow ridge of the medial moraine back towards the station cafe and it was interesting, here, to note the transition from bare rock to closed turf. There were stunning patches of Moss Campion *Silene acaulis*, beautiful deep-blue Spring Gentians and Large-flowered Leopard's-bane *Doronicum grandiflorum*, looking somewhat out of place, among its tiny neighbours.

We enjoyed coffee and cakes in the station cafe, before venturing out again to the start of the Eiger Trail. We descended far enough to see Glacier Crowfoot *Ranunculus glacialis* growing on the cold north-facing screes, before returning past the station to have lunch. The Alpine Choughs provided great entertainment, as they descended to feed on our leftovers. Our afternoon walk took us steeply down to the end of the moraine and then through mountain pasture, vibrant with Globeflower, Alpine Pasqueflower and Red Campion. A broad limestone ridge followed and led us into a forested section before we finally emerged at Wixi. It wasn't long before we were back

in the hotel at Wengernalp, enjoying the delicious home-made apricot tart, I had ordered earlier in the week – a fitting end to a superb week.

Day 8

Sunday 1st July

The good food and hospitality offered by the Hotel Berghaus had, as always, been faultless and we thanked Aldino, Martina and their staff for helping to make such a memorable week. After a relaxed breakfast, we had time for a last look around, before wandering down to the station for our mid-morning train to Zurich.

Receive our e-newsletter

Join the Naturetrek e-mailing list and be the first to hear about new tours, additional departures and new dates, tour reports and special offers. Visit www.naturetrek.co.uk to sign up.

Social Media

We're social! Follow us on Facebook, Twitter and Instagram and be the first to hear about the launch of new tours, offers and exciting sightings and photos from our recently returned holidays.



www.facebook.com/naturetrekwildlifeholidays



www.twitter.com/naturetrektours



www.instagram.com/naturetrek_wildlife_holidays

Species Lists

Plants (‡ = cultivated or naturalised)

| Latin Name | Common Name | Notes |
|----------------------------------|-----------------------|-------------------------------|
| PTERIDOPHYTES | | |
| Aspleniaceae | | |
| <i>Asplenium adiantum-nigrum</i> | Black Spleenwort | Shady cliffs |
| <i>Asplenium ruta-muraria</i> | Wall-rue | Walls |
| <i>Asplenium septentrionale</i> | Forked Spleenwort | Rare, on rocks |
| <i>Asplenium trichomanes</i> | Maidenhair Spleenwort | Walls and cliffs |
| <i>Asplenium viride</i> | Green Spleenwort | Mountain rocks |
| Blechnaceae | | |
| <i>Blechnum spicant</i> | Hard-fern | Locally common |
| Dennstaedtiaceae | | |
| <i>Pteridium aquilinum</i> | Bracken | Valley woodland |
| Dryopteridaceae | | |
| <i>Dryopteris affinis</i> | Scaly Male-fern | Widespread |
| <i>Dryopteris carthusiana</i> | Narrow Buckler-fern | Damp scrub |
| <i>Dryopteris dilatata</i> | Broad Buckler-fern | Frequent |
| <i>Dryopteris expansa</i> | Northern Buckler-fern | Damp scrub |
| <i>Dryopteris filix-mas</i> | Male-fern | Widespread |
| <i>Dryopteris submontana</i> | Rigid Buckler-fern | Limestone rocks. Oberberghorn |
| <i>Gymnocarpium dryopteris</i> | Oak Fern | Woods and rocks. Local |
| <i>Gymnocarpium robertianum</i> | Limestone Fern | Limestone rocks. Oberberghorn |
| <i>Polystichum lonchitis</i> | Holly-fern | Mountain rocks |
| Equisetaceae | | |
| <i>Equisetum arvense</i> | Field Horsetail | Meadows and tracksides |
| <i>Equisetum palustre</i> | Marsh Horsetail | Wet pasture |
| <i>Equisetum sylvaticum</i> | Wood Horsetail | Woods and damp pasture |
| Lycopodiaceae | | |
| <i>Diphasiastrum alpinum</i> | Alpine Clubmoss | Occasional. Mountain turf |
| <i>Huperzia selago</i> | Fir Clubmoss | Frequent. Acidic pasture |
| <i>Lycopodium annotinum</i> | Interrupted Clubmoss | Local. Almend |
| <i>Lycopodium clavatum</i> | Stag's-horn Clubmoss | Local. Above Grutschalp |
| Ophioglossaceae | | |
| <i>Botrychium lunaria</i> | Moonwort | Short turf. Widespread |
| Polypodiaceae | | |
| <i>Polypodium vulgare</i> | Common Polypody | Occasional on rocks |
| Selaginellaceae | | |
| <i>Selaginella selaginoides</i> | Lesser Clubmoss | Occasional in turf |
| Thelypteridaceae | | |
| <i>Oreopteris limbosperma</i> | Lemon-scented Fern | Green Alder scrub and woods |
| <i>Phegopteris connectilis</i> | Beech Fern | Woods. Occasional |

| Latin Name | Common Name | Notes |
|--|--------------------------|------------------------------|
| Woodsiaceae | | |
| <i>Athyrium distentifolium</i> | Lady-fern Family | Mountain woods. Above Murren |
| <i>Athyrium filix-femina</i> | Alpine Lady-fern | Occasional |
| <i>Cystopteris alpina</i> | Lady-fern | Rare. Grutschalp |
| <i>Cystopteris fragilis</i> | Alpine Bladder-fern | Frequent on rocks |
| <i>Cystopteris montana</i> | Brittle Bladder-fern | Local. Trummelbach |
| <i>Woodsia alpina</i> | Mountain Bladder-fern | Rare. Almendhubel |
| <i>Woodsia alpina</i> | Alpine Woodsia | |
| GYMNOSPERMS | | |
| Cupressaceae | | |
| <i>Juniperus communis</i> subsp. <i>nana</i> | Juniper Family | Mountain pasture |
| | Common Juniper | |
| Pinaceae | | |
| <i>Larix decidua</i> | Pine Family | Occasional |
| <i>Picea abies</i> | Larch | Forests |
| <i>Pinus cembra</i> ‡ | Norway Spruce | Occasional |
| <i>Pinus mugo</i> | Arolla Pine | Tree-line |
| <i>Pinus sylvestris</i> | Dwarf Mountain-pine | Widespread |
| <i>Pseudotsuga menziesii</i> ‡ | Scots Pine | Forest below Wengen |
| | Douglas Fir | |
| ANGIOSPERMS | | |
| DICOTYLEDONS | | |
| Adoxaceae | | |
| <i>Adoxa moschatellina</i> | Moschatel Family | Rare. Mannlichen |
| <i>Sambucus ebulus</i> | Moschatel | Lowland. Wilderswil |
| <i>Sambucus nigra</i> | Dwarf Elder | Woods |
| <i>Sambucus racemosa</i> | Common Elder | Occasional. Rocky places |
| <i>Viburnum lantana</i> | Alpine Elder | Woods below Wengen |
| <i>Viburnum opulus</i> | Wayfaring-tree | Woods below Wengen |
| | Guelder-rose | |
| Amaranthaceae | | |
| <i>Chenopodium bonus-henricus</i> | Goosefoot Family | Frequent weed |
| | Good-King-Henry | |
| Apiaceae | | |
| <i>Astrantia major</i> | Carrot Family | Meadows |
| <i>Astrantia minor</i> | Astrantia | Locally common |
| <i>Athamanta cretensis</i> | Lesser Astrantia | Rocky places and moraines |
| <i>Bupleurum stellatum</i> | Athamanta | Mountain rocks |
| <i>Chaerophyllum hirsutum</i> | Starry Hare's-ear | Woods |
| <i>Daucus carota</i> | Hairy Chervil | Wilderswil |
| <i>Heracleum sphondylium</i> | Wild Carrot | Common |
| <i>Imperatoria ostruthium</i> | Hogweed | Common |
| <i>Laserpitium halleri</i> | Masterwort | Occasional |
| <i>Laserpitium latifolium</i> | Haller's Sermountain | Trummelbach |
| <i>Ligusticum mutellina</i> | Broad-leaved Sermountain | Mountain pasture |
| <i>Ligusticum mutellinoides</i> | Alpine Lovage | High-mountain pasture |
| <i>Pimpinella major</i> | Unbranched Lovage | Common |
| <i>Pimpinella saxifraga</i> | Greater Burnet-saxifrage | Meadows |
| | Burnet-saxifrage | |

| Latin Name | Common Name | Notes |
|---|-------------------------------|------------------------------|
| Apocynaceae | Dogbane Family | |
| <i>Vincetoxicum hirundinaria</i> | Swallow-wort | Frequent |
| Araliaceae | Ivy Family | |
| <i>Hedera helix</i> | Ivy | Occasional |
| Asteraceae | Daisy Family | |
| <i>Adenostyles alliariae</i> | Adenostyles | Green Alder scrub |
| <i>Adenostyles glabra</i> | Hairless Adenostyles | Screes. Oberberghorn |
| <i>Adenostyles leucophylla</i> | White-leaved Adenostyles | Screes. Oberberghorn |
| <i>Achillea atrata</i> | Dark-stemmed Sneezewort | Frequent on limestone |
| <i>Achillea erba-rotta</i> subsp. <i>moschata</i> | Musk Milfoil | Acidic mountain pasture |
| <i>Achillea millefolium</i> | Yarrow | Meadows |
| <i>Antennaria carpathica</i> | Carpathian Cat's-foot | Occasional. Mountain pasture |
| <i>Antennaria dioica</i> | Cat's-foot | Common |
| <i>Arnica montana</i> | Arnica | Common |
| <i>Aster alpinus</i> | Alpine Aster | Mountain rocks and turf |
| <i>Bellidastrum michelii</i> | False Aster | Common |
| <i>Bellis perennis</i> | Daisy | Frequent |
| <i>Carduus defloratus</i> | Alpine Thistle | Common |
| <i>Carlina acaulis</i> | Stemless Carline Thistle | Mountain pasture |
| <i>Centaurea montana</i> | Mountain Cornflower | Meadows |
| <i>Centaurea scabiosa</i> | Greater Knapweed | Meadows |
| <i>Cicerbita alpina</i> | Alpine Blue-sowthistle | Forests and scrub |
| <i>Cirsium acaule</i> | Dwarf Thistle | Mountain pasture |
| <i>Cirsium oleraceum</i> | Cabbage Thistle | Common below Wengen |
| <i>Cirsium palustre</i> | Marsh Thistle | Occasional. Wet places |
| <i>Cirsium spinosissimum</i> | Spiniest Thistle | Mountain pasture |
| <i>Crepis aurea</i> | Golden Hawk's-beard | Meadows |
| <i>Crepis paludosa</i> | Marsh Hawk's-beard | Wet meadows |
| <i>Doronicum grandiflorum</i> | Large-flowered Leopard's-bane | Screes |
| <i>Erigeron alpinus</i> | Alpine Fleabane | Mountain rocks. Local |
| <i>Erigeron annuus</i> | Annual Fleabane | Wilderswil |
| <i>Erigeron neglectus</i> | Neglected Fleabane | Mountain rocks. Local |
| <i>Eupatorium cannabinum</i> | Hemp Agrimony | Lowland |
| <i>Gnaphthium supinum</i> | Dwarf Cudweed | Snowbeds |
| <i>Hieracium aurantiacum</i> † | Orange Hawkweed | Wengen |
| <i>Hieracium lactucella</i> | A Hawkweed | Common |
| <i>Hieracium murorum</i> | A Hawkweed | Occasional |
| <i>Hieracium pilosella</i> | Mouse-ear Hawkweed | Common |
| <i>Hieracium villosum</i> | Hairy Hawkweed | Mannlichen |
| <i>Homogyne alpina</i> | Alpine Colt's-foot | Mountain turf. Common |
| <i>Hypochaeris uniflora</i> | Giant Cat's-ear | Mannlichen |
| <i>Leontodon helveticus</i> | Swiss Hawkbit | Mountain turf. Common |
| <i>Leontodon hispidus</i> | Alpine Hawkbit | Acidic pasture |
| <i>Leucanthemopsis alpina</i> | Alpine Moon Daisy | Acidic mountain turf |
| <i>Leucanthemum adustum</i> | Mountain Ox-eye | Mountain pasture |
| <i>Leucanthemum vulgare</i> | Ox-eye Daisy | Meadows |
| <i>Mycelis muralis</i> | Wall Lettuce | Walls |
| <i>Petasites albus</i> | White Butterbur | Mannlichen |
| <i>Petasites paradoxus</i> | Alpine Butterbur | Mountain pasture |
| <i>Prenanthes purpurea</i> | Purple Lettuce | Common. Forests |
| <i>Saussurea alpina</i> | Alpine Saw-wort | Mountain turf |

| Latin Name | Common Name | Notes |
|--|--------------------------------|----------------------------|
| <i>Senecio doronicum</i> | Chamois Ragwort | Limestone rocks and scree |
| <i>Solidago virgaurea</i> | Goldenrod | Mountain pasture |
| <i>Taraxacum alpinum</i> | Alpine Dandelion | Screes |
| <i>Taraxacum officinale</i> | Dandelion | Widespread |
| <i>Tragopogon pratensis</i> | Goat's-beard | Meadows |
| <i>Tussilago farfara</i> | Colt's-foot | Damp places. Widespread |
| Balsaminaceae | Balsam Family | |
| <i>Impatiens parviflora</i> † | Small Balsam | Woods below Wengen |
| Betulaceae | Birch Family | |
| <i>Alnus glutinosa</i> | Common Alder | Valleys, by rivers |
| <i>Alnus incana</i> | Grey Alder | Valleys, by rivers |
| <i>Alnus viridis</i> | Green Alder | Wet mountain slopes |
| <i>Betula pendula</i> | Silver Birch | Widespread |
| <i>Betula pubescens</i> | Downy Birch | Frequent |
| Boraginaceae | Borage Family | |
| <i>Echium vulgare</i> | Viper's Bugloss | Below Wengen |
| <i>Myosotis alpestris</i> | Alpine Forget-me-not | Common |
| Brassicaceae | Cabbage Family | |
| <i>Arabis alpina</i> subsp. <i>alpina</i> | Alpine Rock-cress | Alpine rocks |
| <i>Barbarea vulgaris</i> | Winter-cress | Wet places, mostly lowland |
| <i>Biscutella laevigata</i> | Buckler Mustard | Mountain turf and moraine |
| <i>Capsella bursa-pastoris</i> | Shepherd's-purse | Widespread |
| <i>Cardamine amara</i> | Large Bitter-cress | Streamsides |
| <i>Cardamine flexuosa</i> | Wavy Bitter-cress | Wet places |
| <i>Cardamine hirsuta</i> | Hairy Bitter-cress | Occasional |
| <i>Cardamine impatiens</i> | Narrow-leaved Bitter-cress | Damp places |
| <i>Cardamine pratensis</i> subsp. <i>pratensis</i> | Lady's Smock | Wet places |
| <i>Cardamine resedifolia</i> | Mignonette-leaved Bitter-cress | Mountain rocks |
| <i>Draba aizoides</i> | Yellow Whitlowgrass | Moraines |
| <i>Draba dubia</i> | Austrian Whitlowgrass | Rocky places |
| <i>Draba tomentosa</i> | Wooly Whitlowgrass | Rocks. Almendhubel |
| <i>Erophila verna</i> | Spring Whitlowgrass | Occasional |
| <i>Hornungia alpina</i> subsp. <i>alpina</i> | Chamois Cress | Mountain rocks and screes |
| <i>Kernera saxatilis</i> | Rock Kerneria | Limestone rocks. |
| <i>Sisymbrium austriacum</i> | Austrian Rocket | Waste places |
| <i>Thlaspi rotundifolium</i> subsp. <i>rotundifolium</i> | Round-leaved Penny-cress | Moraines and screes |
| Campanulaceae | Bellflower Family | |
| <i>Campanula barbata</i> | Bearded Bellflower | Mountain turf |
| <i>Campanula cochlearifolia</i> | Fairy's-thimble | Screes |
| <i>Campanula rapunculoides</i> | Creeping Bellflower | Wilderswil |
| <i>Campanula rhomboidalis</i> | Broad-leaved Bellflower | Common in meadows |
| <i>Campanula scheuchzeri</i> | Scheuchzer's Bellflower | Mountain turf |
| <i>Campanula thyrsoidea</i> | Yellow Bellflower | Schynige Platte |
| <i>Campanula trachelium</i> | Nettle-leaved Bellflower | Woods |
| <i>Phyteuma betonicifolium</i> | Betony-leaved Rampion | Acidic pasture |
| <i>Phyteuma hemisphaericum</i> | Globe-headed Rampion | Mountain Rocks |
| <i>Phyteuma orbiculare</i> | Round-headed Rampion | Meadows. Frequent |
| <i>Phyteuma spicatum</i> | Spiked Rampion | Meadows. Frequent |

| Latin Name | Common Name | Notes |
|--|--------------------------------|----------------------------|
| Cannabaceae | Hop Family | |
| <i>Humulus lupulus</i> | Hop | Below Wengen |
| Caprifoliaceae | Honeysuckle Family | |
| <i>Knautia dipsacifolia</i> | Wood Scabious | Common |
| <i>Lonicera alpigena</i> | Alpine Honeysuckle | Occasional. Forests |
| <i>Lonicera caerulea</i> | Blue-berried Honeysuckle | Green Alder scrub |
| <i>Lonicera xylostemum</i> | Fly Honeysuckle | Woods |
| <i>Scabiosa columbaria</i> | Small Scabious | Common in lowlands |
| <i>Scabiosa lucida</i> | Shining Scabious | Meadows |
| <i>Succisa pratensis</i> | Devil's-bit Scabious | Damp meadows |
| <i>Valeriana dioica</i> | Marsh Valerian | Wet pasture |
| <i>Valeriana officinalis</i> | Common Valerian | Damp meadows |
| <i>Valeriana tripteris</i> | Three-leaved Valerian | Woods |
| Caryophyllaceae | Pink Family | |
| <i>Arenaria biflora</i> | Two-flowered Sandwort | Screens |
| <i>Cerastium arvense</i> subsp. <i>strictum</i> | Field Mouse-ear | Frequent |
| <i>Cerastium cerastoides</i> | Starwort Mouse-ear | Rocks and pasture |
| <i>Cerastium fontanum</i> | Common Mouse-ear | Common |
| <i>Cerastium latifolium</i> | Broad-leaved Mouse-ear | Moraines and screes |
| <i>Dianthus carthusianorum</i> | Carthusian Pink | Almendhubel |
| <i>Dianthus superbus</i> ‡ | Large Pink | Railway banks below Almond |
| <i>Dianthus sylvestris</i> | Wood Pink | Rocks. Local |
| <i>Gypsophila repens</i> | Alpine Gypsophila | Frequent |
| <i>Minuartia sedoides</i> | Mossy Cyphel | Alpine turf |
| <i>Minuartia verna</i> | Vernal Sandwort | Rocky places |
| <i>Moehringia ciliata</i> | Creeping Sandwort | Screens |
| <i>Moehringia muscosa</i> | Mossy Sandwort | Shady places. Local |
| <i>Sagina procumbens</i> | Procumbent Pearlwort | Common |
| <i>Sagina saginoides</i> | Alpine Pearlwort | Alpine turf and screes |
| <i>Silene acaulis</i> | Moss Campion | Mountain turf |
| <i>Silene dioica</i> | Red Campion | Frequent. Meadows |
| <i>Silene nutans</i> | Nottingham Catchfly | Occasional. Rocks |
| <i>Silene rupestris</i> | Rock Campion | Common on rocks |
| <i>Silene vulgaris</i> | Bladder Campion | Occasional. Meadows |
| <i>Stellaria graminea</i> | Lesser Stitchwort | Occasional |
| <i>Stellaria media</i> | Common Chickweed | Occasional |
| <i>Stellaria nemorum</i> | Wood Stitchwort | Woods |
| Celastraceae | Spindle Family | |
| <i>Parnassia palustris</i> | Grass-of-Parnassus | Damp flushes |
| Cistaceae | Rockrose Family | |
| <i>Helianthemum nummularium</i> subsp. <i>grandiflorum</i> | Large-flowered Common Rockrose | Common |
| <i>Helianthemum oelandicum</i> subsp. <i>alpestre</i> | Alpine Rockrose | Frequent |
| Cornaceae | Dogwood Family | |
| <i>Cornus sanguinea</i> | Dogwood | Woods |
| Corylaceae | Hazel Family | |
| <i>Corylus avellana</i> | Hazel | Woods |

| Latin Name | Common Name | Notes |
|---|----------------------------|---------------------------|
| Crassulaceae | | |
| Stonecrop Family | | |
| <i>Rhodiola rosea</i> ‡ | Roseroot | Alpine Garden |
| <i>Sedum acre</i> | Biting Stonecrop | Occasional |
| <i>Sedum album</i> | White Stonecrop | Walls. Wengen |
| <i>Sedum alpestre</i> | Alpine Stonecrop | Mountain rocks |
| <i>Sedum atratum</i> | Dark Stonecrop | Mountain rocks |
| <i>Sedum dasyphyllum</i> | Thick-leaved Stonecrop | Walls and rocks |
| <i>Sedum hispanicum</i> ‡ | Spanish Stonecrop | Naturalised. Wengen |
| <i>Sedum spurium</i> ‡ | Caucasian Stonecrop | Wengen |
| <i>Sedum rubens</i> | Reddish Stonecrop | Rocks |
| <i>Sempervivum montanum</i> | Mountain Houseleek | Common on rocks |
| Cuscutaceae | | |
| Dodder Family | | |
| <i>Cuscuta epithymum</i> | Common Dodder | Frequent |
| Ericaceae | | |
| Heather Family | | |
| <i>Arctostaphylos alpinus</i> | Alpine Bearberry | Occasional. Mountain turf |
| <i>Arctostaphylos uva-ursi</i> | Bearberry | Acidic heaths |
| <i>Calluna vulgaris</i> | Ling | Frequent |
| <i>Empetrum nigrum</i> subsp. <i>hermaphroditum</i> | Hermaphrodite Crowberry | Acidic heaths |
| <i>Erica carnea</i> | Spring Heath | Frequent on limestone |
| <i>Kalmia procumbens</i> | Trailing Azalea | Acidic heaths |
| <i>Moneses uniflora</i> | One-flowered Wintergreen | Forest |
| <i>Orthilia secunda</i> | Serrated Wintergreen | Forest |
| <i>Pyrola minor</i> | Lesser Wintergreen | Forest |
| <i>Rhododendron ferrugineum</i> | Alpenrose | Common |
| <i>Rhododendron hirsutum</i> | Hairy Alpenrose | Common on limestone |
| <i>Vaccinium myrtillus</i> | Bilberry | Common |
| <i>Vaccinium uliginosum</i> agg. | Bog Whortleberry | Common |
| <i>Vaccinium vitis-idaea</i> | Cowberry | Common |
| Euphorbiaceae | | |
| Spurge Family | | |
| <i>Euphorbia cyparissias</i> | Cypress Spurge | Frequent on limestone |
| <i>Mercurialis perennis</i> | Dog's Mercury | Woods |
| Fabaceae | | |
| Pea Family | | |
| <i>Anthyllis vulneraria</i> subsp. <i>alpestris</i> | Alpine Kidney Vetch | Mountain pasture |
| <i>Astragalus alpinus</i> | Alpine Milk-vetch | Mountain turf |
| <i>Astragalus australis</i> | Southern Milk-vetch | Mountain turf |
| <i>Astragalus frigidus</i> | Pallid Milk-vetch | Limestone cliffs |
| <i>Coronilla emerus</i> | False Senna | Occasional |
| <i>Coronilla vaginalis</i> | Small Scorpion-vetch | Cliffs below Wengen |
| <i>Hedysarum hedysaroides</i> | Alpine Sainfoin | Mountain pasture |
| <i>Hippocrepis comosa</i> | Horseshoe Vetch | Common |
| <i>Lathyrus laevigatus</i> subsp. <i>occidentalis</i> | Yellow Pea | Oberberghorn |
| <i>Lathyrus pratensis</i> | Meadow Vetchling | Meadows |
| <i>Lotus alpinus</i> | Alpine Bird's-foot-trefoil | Mountain turf |
| <i>Melilotus albus</i> | White Melilot | Lowland |
| <i>Onobrychis arenaria</i> | Small Sainfoin | Occasional |
| <i>Onobrychis montana</i> | Mountain Sainfoin | Mountain pasture |
| <i>Ononis repens</i> | Common Restharrow | Below Wengen |
| <i>Oxytropis campestris</i> | Yellow Milk-vetch | Locally common |
| <i>Oxytropis jacquinii</i> | Mountain Milk-vetch | Moraines. Local |

| Latin Name | Common Name | Notes |
|-------------------------------|------------------------------|-------------------------------|
| <i>Robinia pseudoacacia</i> † | False Acacia | Lowlands |
| <i>Securigera varia</i> | Crown Vetch | Wilderswil |
| <i>Trifolium alpinum</i> | Alpine Clover | Common |
| <i>Trifolium badium</i> | Brown Clover | Mountain meadows |
| <i>Trifolium montanum</i> | Mountain Clover | Mountain pasture |
| <i>Trifolium pratense</i> | Red Clover | Meadows |
| <i>Trifolium repens</i> | White Clover | Frequent |
| <i>Trifolium thalii</i> | Thal's Clover | Mountain turf and scree |
| <i>Vicia cracca</i> | Tufted Vetch | Frequent |
| <i>Vicia sepium</i> | Bush Vetch | Occasional |
| <i>Vicia sylvatica</i> | Wood Vetch | Woods |
| Fagaceae | Beech Family | |
| <i>Castanea sativa</i> † | Sweet Chestnut | Occasional |
| <i>Fagus sylvatica</i> | Beech | Valley forest |
| <i>Quercus petraea</i> | Sessile Oak | Frequent. Woods |
| <i>Quercus robur</i> | Pedunculate Oak | Occasional |
| Gentianaceae | Gentian Family | |
| <i>Gentiana acaulis</i> | Trumpet Gentian | Mountain pasture |
| <i>Gentiana asclepiadea</i> | Willow Gentian | Woods on limestone |
| <i>Gentiana bavarica</i> | Bavarian Gentian | Marshes |
| <i>Gentiana brachyphylla</i> | Short-leaved Gentian | Alpine turf |
| <i>Gentiana clusii</i> | Clusius' Gentian | Limestone pasture |
| <i>Gentiana lutea</i> | Great Yellow Gentian | Mountain pasture |
| <i>Gentiana purpurea</i> | Purple Gentian | Green Alder scrub and pasture |
| <i>Gentiana verna</i> | Spring Gentian | Mountain pasture |
| Geraniaceae | Crane's-bill Family | |
| <i>Geranium pyrenaicum</i> | Pyrenean Crane's-bill | Occasional |
| <i>Geranium robertianum</i> | Herb Robert | Occasional |
| <i>Geranium sylvaticum</i> | Wood Crane's-bill | Common |
| Hypericaceae | St John's-wort Family | |
| <i>Hypericum maculatum</i> | Imperforate St John's-wort | Frequent |
| Juglandaceae | Walnut Family | |
| <i>Juglans regia</i> † | Walnut | Occasional |
| Lamiaceae | Dead-nettle Family | |
| <i>Ajuga pyramidalis</i> | Pyramidal Bugle | Locally common |
| <i>Ajuga reptans</i> | Common Bugle | Common |
| <i>Betonica officinalis</i> | Betony | Meadows |
| <i>Clinopodium alpinum</i> | Alpine Calamint | Occasional. Rocky banks |
| <i>Clinopodium vulgare</i> | Wild Basil | Frequent |
| <i>Galeopsis tetrahit</i> | Common Hemp-nettle | Occasional |
| <i>Glechoma hederacea</i> | Ground-ivy | Woods |
| <i>Lamium galeobdolon</i> | Yellow Archangel | Woods |
| <i>Lamium maculatum</i> | Spotted Dead-nettle | Occasional |
| <i>Lamium purpureum</i> | Red Dead-nettle | Occasional |
| <i>Origanum vulgare</i> | Marjoram | Occasional. Meadows |
| <i>Prunella grandiflora</i> | Large-flowered Self-heal | Meadows. Trummelbach |
| <i>Prunella vulgaris</i> | Self-heal | Common |

| Latin Name | Common Name | Notes |
|--|--------------------------------------|------------------------------|
| <i>Salvia glutinosa</i> | Sticky Clary | Forest. Wilderwil |
| <i>Salvia pratensis</i> | Meadow Clary | Meadows. Trummelbach |
| <i>Stachys alpina</i> | Alpine Woundwort | Below Wengen |
| <i>Stachys sylvatica</i> | Hedge Woundwort | Woods |
| <i>Teucrium chamaedrys</i> | Wall Germander | Below Wengen |
| <i>Teucrium montanum</i> | Mountain Germander | Trummelbach |
| <i>Teucrium scorodonia</i> | Wood Sage | Woods |
| <i>Thymus polytrichus</i> | Wild Thyme | Common |
| Lentibulariaceae | Butterwort Family | |
| <i>Pinguicula alpina</i> | Alpine Butterwort | Mountain rocks and turf |
| <i>Pinguicula vulgaris</i> | Common Butterwort | Wet turf |
| Linaceae | Flax Family | |
| <i>Linum alpinum</i> † | Alpine Flax | Wengernalp and Alpine Garden |
| <i>Linum catharticum</i> | Fairy Flax | Common |
| Oleaceae | Ash Family | |
| <i>Fraxinus excelsior</i> | Ash | Lowland |
| <i>Ligustrum vulgare</i> | Wild Privet | Lowland |
| Onagraceae | Willowherb Family | |
| <i>Chamerion angustifolium</i> | Rosebay Willowherb | Occasional |
| <i>Epilobium alpestre</i> | Whorled-leaved Willowherb | Frequent |
| <i>Epilobium alsinifolium</i> | Chickweed Willowherb | Frequent |
| <i>Epilobium anagallidifolium</i> | Pimpernel-leaved Willowherb | Frequent |
| <i>Epilobium hirsutum</i> | Great Willowherb | Lowlands |
| <i>Epilobium montanum</i> | Broad-leaved Willowherb | Occasional |
| Orobanchaceae | Broomrape Family | |
| <i>Bartsia alpina</i> | Alpine Bartsia | Mountain turf |
| <i>Euphrasia minima</i> | Dwarf Eyebright | Local. Mountain pasture |
| <i>Euphrasia officinalis</i> subsp. <i>monticola</i> | Montane Eyebright | Mountain pasture |
| <i>Euphrasia officinalis</i> subsp. <i>pratensis</i> | Rostkov's Eyebright | Valley meadows |
| <i>Melampyrum pratense</i> | Common Cow-wheat | Woods |
| <i>Melampyrum sylvaticum</i> | Small Cow-wheat | Forest |
| <i>Orobanche alba</i> | Thyme Broomrape | Occasional |
| <i>Orobanche caryophyllacea</i> | Clove-scented Broomrape | Occasional |
| <i>Pedicularis foliosa</i> | Leafy Lousewort | Local. Limestone pasture |
| <i>Pedicularis oederi</i> | Crimson-tipped Lousewort | Alpine turf |
| <i>Pedicularis tuberosa</i> | Long-beaked Yellow Lousewort | Acidic turf |
| <i>Pedicularis verticillata</i> | Whorled Lousewort | Mountain turf |
| <i>Rhinanthus alectorolophus</i> | Greater Yellow-rattle | Common. Meadows |
| <i>Rhinanthus glacialis</i> | Glacier Yellow-rattle | Mountain pasture |
| <i>Rhinanthus minor</i> | Yellow-rattle | Uncommon. Pasture |
| Oxalidaceae | Wood-sorrel Family | |
| <i>Oxalis acetosella</i> | Wood-sorrel | Shaded rocks |
| Plantaginaceae | Plantain and Speedwell Family | |
| <i>Digitalis grandiflora</i> | Large Yellow Foxglove | Local. Forest |
| <i>Erinus alpinus</i> | Fairy Foxglove | Frequent |
| <i>Globularia cordifolia</i> | Matted Globularia | Frequent |

| Latin Name | Common Name | Notes |
|---|-----------------------------|----------------------------|
| <i>Globularia nudicaulis</i> | Leafless-stemmed Globularia | Frequent |
| <i>Linaria alpina</i> | Alpine Toadflax | Moraine and scree |
| <i>Plantago alpina</i> | Alpine Plantain | Mountain turf |
| <i>Plantago atrata</i> | Dark Plantain | Mountain turf |
| <i>Plantago major</i> | Greater Plantain | Occasional |
| <i>Plantago maritima</i> subsp. <i>serpentina</i> | Fleshy Plantain | Frequent |
| <i>Plantago media</i> | Hoary Plantain | Occasional |
| <i>Plantago lanceolata</i> | Ribwort Plantain | Occasional |
| <i>Veronica alpina</i> | Alpine Speedwell | Stony mountain pasture |
| <i>Veronica aphylla</i> | Leafless-stemmed Speedwell | Stony mountain pasture |
| <i>Veronica beccabunga</i> | Brooklime | Wet places |
| <i>Veronica bellidioides</i> | Violet Speedwell | Acidic pasture |
| <i>Veronica chamaedrys</i> | Germander Speedwell | Below Wengen |
| <i>Veronica fruticans</i> | Rock Speedwell | Mountain rocks |
| <i>Veronica fruticososa</i> | A Rock Speedwell | Oberberghorn |
| <i>Veronica officinalis</i> | Heath Speedwell | Woods |
| <i>Veronica serpyllifolia</i> | Thyme-leaved Speedwell | Frequent |
| <i>Veronica urticifolia</i> | Nettle-leaved Speedwell | Woods |
| Polygalaceae | Milkwort Family | |
| <i>Polygala alpestris</i> | Mountain Milkwort | Common |
| <i>Polygala chamaebuxus</i> | Shrubby Milkwort | Frequent on rocks |
| <i>Polygala vulgaris</i> | Common Milkwort | Frequent |
| Polygonaceae | Dock Family | |
| <i>Fallopia japonica</i> ‡ | Japanese Knotweed | Lowland |
| <i>Oxyria digyna</i> | Mountain Sorrel | Mountain rocks and scree |
| <i>Persicaria bistorta</i> | Common Bistort | Damp pasture |
| <i>Persicaria vivipara</i> | Alpine Bistort | Common. Mountain turf |
| <i>Rumex acetosa</i> | Common Sorrel | Widespread |
| <i>Rumex acetosella</i> | Sheep's Sorrel | Acidic pasture |
| <i>Rumex alpestris</i> | Mountain Dock | Mountain pasture |
| <i>Rumex alpinus</i> | Monk's-rhubarb | Disturbed mountain pasture |
| <i>Rumex obtusifolius</i> | Broad-leaved Dock | Occasional |
| <i>Rumex scutatus</i> | French Sorrel | Rocks and scree |
| Primulaceae | Primrose Family | |
| <i>Androsace chamaejasme</i> | Ciliate Rock-jasmine | Frequent |
| <i>Androsace helvetica</i> | Swiss Rock-jasmine | Alpine cliffs |
| <i>Androsace obtusifolia</i> | Blunt-leaved Rock-jasmine | Acidic turf |
| <i>Androsace pubescens</i> | Hairy Rock-jasmine | Rare. Mountain rocks |
| <i>Lysimachia nemorum</i> | Yellow Pimpernel | Damp woods |
| <i>Primula auricula</i> | Auricula | Mountain rocks and turf |
| <i>Primula elatior</i> | Oxlip | Snowbeds |
| <i>Primula farinosa</i> | Bird's-eye Primrose | Mountain turf |
| <i>Primula hirsuta</i> | Hairy Primrose | Acidic rocks |
| <i>Primula veris</i> | Cowslip | Below Wengen |
| <i>Soldanella alpina</i> | Alpine Snowbell | Common |
| Ranunculaceae | Buttercup Family | |
| <i>Aconitum lycoctonum</i> subsp. <i>vulparia</i> | Wolf's-bane | Frequent. Forest |
| <i>Aconitum napellus</i> agg. | Monk's-hood | Damp meadows |
| <i>Actaea spicata</i> | Baneberry | Woods |

| Latin Name | Common Name | Notes |
|--|----------------------------|-------------------------------|
| <i>Anemone narcissiflora</i> | Narcissus-flowered Anemone | Mountain pasture |
| <i>Anemone nemorosa</i> | Wood Anemone | Woods |
| <i>Aquilegia alpina</i> | Alpine Columbine | Rare. Biglenalp, Eiger |
| <i>Aquilegia atrata</i> | Dark Columbine | Forest and meadows |
| <i>Caltha palustris</i> | Marsh-marigold | Flushes and streamsides |
| <i>Clematis alpina</i> ‡ | Alpine Clematis | Alpine Garden |
| <i>Clematis vitalba</i> | Old Man's Beard | Common. Lowland |
| <i>Hepatica nobilis</i> | Liverleaf | Woodland |
| <i>Pulsatilla alpina</i> subsp. <i>alpina</i> | Alpine Pasqueflower | Frequent on limestone |
| <i>Pulsatilla alpina</i> subsp. <i>apiifolia</i> | Alpine Pasqueflower | Frequent on acidic soils |
| <i>Pulsatilla vernalis</i> | Spring Pasqueflower | Locally common |
| <i>Ranunculus aconitifolius</i> | Aconite-leaved Buttercup | Locally common |
| <i>Ranunculus acris</i> | Meadow Buttercup | Common |
| <i>Ranunculus alpestris</i> | Alpine Buttercup | Snowbeds. Common |
| <i>Ranunculus glacialis</i> | Glacier Crowfoot | Alpine screes. Eiger |
| <i>Ranunculus kuepferi</i> | Kupfer's Buttercup | Locally common |
| <i>Ranunculus montanus</i> | Mountain Buttercup | Mountain pasture |
| <i>Ranunculus parnassifolius</i> ‡ | Parnassus-leaved Buttercup | Alpine Garden |
| <i>Ranunculus repens</i> | Creeping Buttercup | Occasional. Meadows |
| <i>Thalictrum alpinum</i> | Alpine Meadow-rue | Uncommon. Alpine turf |
| <i>Thalictrum aquilegifolium</i> | Great Meadow-rue | Mountain meadows |
| <i>Thalictrum foetidum</i> | Stinking Meadow-rue | Trummelbach |
| <i>Trollius europaeus</i> | Globeflower | Common |
| Rhamnaceae | Buckthorn Family | |
| <i>Rhamnus alpinus</i> | Alpine Buchthorn | Rocky places |
| <i>Rhamnus pumilus</i> | Dwarf Buckthorn | Limestone rocks. Oberberghorn |
| Rosaceae | Rose Family | |
| <i>Agrimonia eupatoria</i> | Agrimony | Occasional. Lowland |
| <i>Alchemilla alpina</i> | Alpine Lady's-mantle | Common |
| <i>Alchemilla conjuncta</i> | Silver Lady's-mantle | Frequent |
| <i>Alchemilla glabra</i> | Smooth Lady's-mantle | Frequent |
| <i>Alchemilla pentaphyllea</i> | Cut-leaved Lady's-mantle | Snowbeds |
| <i>Alchemilla vulgaris</i> agg. | Lady's-mantle | Frequent |
| <i>Alchemilla xanthochlora</i> | Pale Lady's-mantle | Frequent |
| <i>Aruncus dioicus</i> | Goat's-beard Spiraea | Woods. Below Wengen |
| <i>Cotoneaster integerrima</i> | Wild Cotoneaster | Mountain rocks. Oberberghorn |
| <i>Crataegus monogyna</i> | Hawthorn | Woods |
| <i>Dryas octopetala</i> | Mountain Avens | Mountain rocks and scree |
| <i>Filipendula ulmaria</i> | Meadowsweet | Lowland meadows |
| <i>Fragaria vesca</i> | Wild Strawberry | Woods and pasture |
| <i>Geum montanum</i> | Alpine Avens | Mountain turf |
| <i>Geum rivale</i> | Water Avens | Occasional. Wet pasture |
| <i>Geum urbanum</i> | Herb Bennet | Woods |
| <i>Potentilla crantzii</i> | Alpine Cinquefoil | Common. Mountain turf |
| <i>Potentilla erecta</i> | Tormentil | Acidic turf |
| <i>Potentilla sterilis</i> | Barren Strawberry | Frequent |
| <i>Poterium sanguisorba</i> | Salad Burnet | Widespread |
| <i>Prunus avium</i> | Wild Cherry | Woods |
| <i>Rosa canina</i> | Dog-rose | Below Wengen |
| <i>Rosa glauca</i> | Blue-leaved Rose | Occasional |
| <i>Rosa micrantha</i> | Small-flowered Sweet-briar | Occasional |

| Latin Name | Common Name | Notes |
|---|--------------------------------|---------------------------|
| <i>Rosa pendulina</i> | Alpine Rose | Frequent. Mountains |
| <i>Rosa rubiginosa</i> | Sweet-briar | Below Wengen |
| <i>Rubus idaeus</i> | Raspberry | Woods |
| <i>Rubus saxatilis</i> | Stone Bramble | Occasional. Rocky places |
| <i>Sibbaldia procumbens</i> | Sibbaldia | Snowbeds |
| <i>Sorbus aria</i> | Whitebeam | Occasional. Woods |
| <i>Sorbus aucuparia</i> | Rowan | Woods |
| <i>Sorbus chamaemespilus</i> | False-medlar | Mountain pasture |
| Rubiaceae | Madder Family | |
| <i>Galium megalospermum</i> | Swiss Bedstraw | Moraines |
| <i>Galium mollugo</i> | Hedge Bedstraw | Meadows |
| <i>Galium odoratum</i> | Woodruff | Woods |
| <i>Galium pumilum</i> | Slender Bedstraw | Frequent |
| <i>Galium verum</i> | Lady's Bedstraw | Meadows |
| Salicaceae | Willow Family | |
| <i>Populus alba</i> | White Poplar | Lowland |
| <i>Populus tremula</i> | Aspen | Occasional |
| <i>Salix aurita</i> | Eared Willow | Widespread |
| <i>Salix breviserrata</i> | Finely-toothed Willow | Occasional |
| <i>Salix hastata</i> | Large-stipuled Willow | Local. Lauberhorn |
| <i>Salix glaucosericea</i> | Silky Willow | Local. Lauberhorn |
| <i>Salix hegetschweileri</i> | Alpine Willow | Mountain turf |
| <i>Salix herbacea</i> | Least Willow | Mountain turf |
| <i>Salix reticulata</i> | Net-leaved Willow | Alpine rocks and turf |
| <i>Salix retusa</i> | Retuse-leaved Willow | Alpine rocks and turf |
| <i>Salix serpyllifolia</i> | Thyme-leaved Willow | Alpine rocks and turf |
| Santalaceae | Bastard-toadflax Family | |
| <i>Thesium alpinum</i> | Alpine Bastard-toadflax | Occasional. Pasture |
| Sapindaceae | Maple Family | |
| <i>Acer campestre</i> | Field Maple | Frequent |
| <i>Acer platanoides</i> | Norway Maple | Occasional |
| <i>Acer pseudoplatanus</i> | Sycamore | Common |
| Saxifragaceae | Saxifrage Family | |
| <i>Saxifraga aizoides</i> | Yellow Mountain Saxifrage | Wet turf and scree |
| <i>Saxifraga androsacea</i> | Scree Saxifrage | Damp scree and snowbeds |
| <i>Saxifraga bryoides</i> | Mossy Saxifrage | Acidic rocks |
| <i>Saxifraga caesia</i> | Blue Saxifrage | Limestone rocks and scree |
| <i>Saxifraga exarata</i> subsp. <i>moschata</i> | Musky Saxifrage | Mountain rocks |
| <i>Saxifraga oppositifolia</i> | Purple Saxifrage | Rocks, moraine and scree |
| <i>Saxifraga paniculata</i> | Livelong Saxifrage | Rocks |
| <i>Saxifraga rotundifolia</i> | Round-leaved Saxifrage | Damp shaded rocks |
| <i>Saxifraga stellaris</i> | Starry Saxifrage | Acidic flushes |
| Scrophulariaceae | Figwort Family | |
| <i>Verbascum nigrum</i> | Dark Mullein | Lowland |
| Solanaceae | Nightshade Family | |
| <i>Atropa bella-donna</i> | Deadly Nightshade | Forest. Wilderswil |

| Latin Name | Common Name | Notes |
|---|---|--|
| Thymelaeaceae <i>Daphne mezereum</i> | Mezereon Family Mezereon | Local. Mountain pasture |
| Tiliaceae <i>Tilia cordata</i> <i>Tilia platyphyllos</i> | Lime Family Small-leaved Lime Large-leaved Lime | Lowland Lowland |
| Ulmaceae <i>Ulmus glabra</i> | Elm Family Wych Elm | Valley woods |
| Urticaceae <i>Urtica dioica</i> | Nettle Family Common Nettle | Widespread |
| Violaceae <i>Viola biflora</i> <i>Viola calcarata</i> <i>Viola cenisia</i> <i>Viola lutea</i> <i>Viola palustris</i> <i>Viola riviniana</i> <i>Viola tricolor</i> | Violet Family Yellow Wood Violet Long-spurred Pansy Mt Cenis Pansy Mountain Pansy Marsh Violet Common Dog-violet Wild Pansy | Common Locally common Moraine and scree Locally common Common Common Local |
| MONOCOTYLEDONS | | |
| Amaryllidaceae <i>Allium schoenoprasum</i> <i>Allium victorialis</i> | Daffodil Family Chives Alpine Leek | Occasional. Mountains Uncommon. Mountain pasture |
| Asparagaceae <i>Anthericum liliago</i> <i>Anthericum ramosum</i> <i>Convallaria majalis</i> <i>Maianthemum bifolium</i> <i>Paradisea liliastrum</i> <i>Polygonatum multiflorum</i> <i>Polygonatum odoratum</i> <i>Polygonatum verticillatum</i> | Asparagus Family St Bernard's Lily Branched St Bernard's Lily Lily-of-the-Valley May Lily St Bruno's Lily Solomon's-seal Angular Solomon's-seal Whorled Solomon's-seal | Meadow above Wengen Trummelbach Limestone cliffs and scree Mountain woods Local. Mountain pasture Lowland woods Limestone cliffs and scree Occasional. Forests |
| Colchicaceae <i>Colchicum autumnale</i> | Meadow Saffron Family Meadow Saffron | Meadows below Wengen |
| Cyperaceae <i>Carex atrata</i> <i>Carex canescens</i> <i>Carex capillaris</i> <i>Carex caryophyllea</i> <i>Carex curvula</i> <i>Carex davalliana</i> <i>Carex echinata</i> <i>Carex ferruginea</i> <i>Carex flacca</i> <i>Carex flava</i> <i>Caex nigra</i> | Sedge Family Black Alpine Sedge White Sedge Hair Sedge Spring Sedge An alpine Sedge Davall's Sedge Star Sedge Rusty Sedge Glaucous Sedge Yellow Sedge Black Sedge | Wet mountain pasture Wet mountain pasture Wet mountain pasture Dry pasture Dominant. Mountain turf Wet calcareous turf Frequent Dominant. Limestone turf Common Marshes Common |

| Latin Name | Common Name | Notes |
|--|---|---------------------------------|
| <i>Carex panicea</i> | Carnation Sedge | Common |
| <i>Carex rostrata</i> | Bottle Sedge | Pond margins |
| <i>Carex sempervirens</i> | Evergreen Sedge | Mountain grassland on limestone |
| <i>Eriophorum angustifolium</i> | Common Cottongrass | Wet flushes and bogs |
| <i>Eriophorum latifolium</i> | Broad-leaved Cottongrass | Marshes |
| <i>Eriophorum scheuchzeri</i> | Scheuchzer's Cottongrass | Occasional. Bogs |
| <i>Eriophorum vaginatum</i> | Hare's-tail Cottongrass | Acidic bogs |
| Iridaceae | | |
| <i>Crocus vernus</i> | Iris Family Spring Crocus | Common |
| Juncaceae | | |
| <i>Juncus alpinoarticulatus</i> | Rush Family Alpine Rush | Wet mountain pasture |
| <i>Juncus jacquinii</i> | Jacquin's Rush | Wet mountain pasture |
| <i>Juncus trifidus</i> | Three-leaved Rush | Acidic mountain turf |
| <i>Juncus triglumis</i> | Three-flowered Rush | Acidic mountain turf |
| <i>Luzula alpinopilosa</i> | Alpine Wood-rush | Mountain turf |
| <i>Luzula lutea</i> | Yellow Wood-rush | Mountain turf |
| <i>Luzula multiflora</i> | Heath Wood-rush | Mountain turf |
| <i>Luzula spicata</i> | Spiked Wood-rush | Mountain turf |
| <i>Luzula sylvatica</i> | Great Wood-rush | Woods |
| Liliaceae | | |
| <i>Gagea fistulosa</i> | Lily Family A Yellow Star-of-Bethlehem | Uncommon. Mountain turf |
| <i>Gagea serotina</i> | Snowdon Lily | Mountain turf |
| <i>Lilium martagon</i> | Martagon Lily | Woods |
| <i>Streptopus amplexifolius</i> | Streptopus | Occasional. Forests |
| Melanthiaceae | | |
| <i>Paris quadrifolia</i> | Herb-Paris Family Herb-Paris | Local. Woods |
| <i>Veratrum album</i> | White False-helleborine | Mountain pasture |
| Orchidaceae | | |
| <i>Cephalantha longifolia</i> | Orchid Family Narrow-leaved Helleborine | Frequent. Woods |
| <i>Cephalanthera rubra</i> | Red Helleborine | Trummelbach |
| <i>Chamorchis alpina</i> | Dwarf Orchid | Local |
| <i>Coeloglossum viride</i> | Frog Orchid | Frequent |
| <i>Corallorhiza trifida</i> | Coralroot Orchid | Local. Forests |
| <i>Cypripedium calceolus</i> | Lady's-slipper | Local |
| <i>Dactylorhiza fuchsii</i> | Common Spotted-orchid | Damp base-rich soils |
| <i>Dactylorhiza maculata</i> | Heath Spotted-orchid | Acidic soils |
| <i>Dactylorhiza majalis</i> | Broad-leaved Marsh-orchid | Common |
| <i>Epipactis atrorubens</i> | Dark-red Helleborine | Frequent. Forests |
| <i>Epipactis helleborine</i> | Broad-leaved Helleborine | Frequent. Forests |
| <i>Gymnadenia conopsea</i> | Fragrant-orchid | Common |
| <i>Gymnadenia odoratissima</i> | Short-spurred Fragrant-orchid | Trummelbach |
| <i>Gymnadenia rhellicani</i> | Vanilla Orchid | Locally common |
| <i>Gymnadenia x suaveolens</i> (<i>G.conopsea</i> x <i>G.rhellicani</i>) | A hybrid orchid | Wixi |
| <i>Neotinea ustulata</i> | Burnt Orchid | Uncommon. Mountain pasture |
| <i>Neottia cordata</i> | Lesser Twayblade | Locally common |
| <i>Neottia nidus-avis</i> | Bird's-nest Orchid | Frequent. Forests |
| <i>Neottia ovata</i> | Twayblade | Locally common |

| Latin Name | Common Name | Notes |
|--------------------------------|---------------------------------|----------------------------|
| <i>Orchis mascula</i> | Early-purple Orchid | Common |
| <i>Platanthera bifolia</i> | Lesser Butterfly-orchid | Locally frequent. Forests |
| <i>Platanthera chlorantha</i> | Greater Butterfly-orchid | Local. Woods |
| <i>Pseudorchis albida</i> | Small-white Orchid | Common. Mountain turf |
| <i>Traunsteinera globosa</i> | Round-headed Orchid | Local |
| Poaceae | Grass Family | |
| <i>Agropyron repens</i> | Common Couch | Occasional |
| <i>Agrostis capillaris</i> | Common Bent | Common |
| <i>Anthoxanthum alpinum</i> | Alpine Vernal-grass | Frequent. Mountain turf |
| <i>Anthoxanthum odoratum</i> | Sweet Vernal-grass | Common |
| <i>Arrhenatherum elatius</i> | False Oat-grass | Common |
| <i>Brachypodium sylvaticum</i> | False Brome | Woods |
| <i>Briza media</i> | Quaking-grass | Locally common |
| <i>Bromus hordeaceus</i> | Soft Brome | Frequent |
| <i>Bromus sterilis</i> | Barren Brome | Frequent |
| <i>Cynosurus cristatus</i> | Crested Dog's-tail | Common |
| <i>Dactylis glomerata</i> | Cock's-foot | Common |
| <i>Deschampsia cespitosa</i> | Tufted Hair-grass | Wet mountain pasture |
| <i>Deschampsia flexuosa</i> | Wavy Hair-grass | Acidic mountain turf |
| <i>Festuca ovina</i> | Sheep's-fescue | Common |
| <i>Festuca rubra</i> | Red Fescue | Common |
| <i>Festuca violacea</i> | Violet Fescue | Mountain turf |
| <i>Holcus lanatus</i> | Yorkshire-fog | Locally common |
| <i>Holcus mollis</i> | Creeping Soft-grass | Locally common |
| <i>Koeleria macrantha</i> | Crested Hair-grass | Frequent |
| <i>Lolium perenne</i> | Perennial Rye-grass | Frequent |
| <i>Melica uniflora</i> | Wood Melick | Woods |
| <i>Nardus stricta</i> | Mat-grass | Common |
| <i>Phleum alpinum</i> | Alpine Timothy | Frequent. Mountain pasture |
| <i>Phleum pratense</i> | Timothy | Common |
| <i>Poa alpina</i> | Alpine Meadow-grass | Common |
| <i>Poa annua</i> | Annual Meadow-grass | Common on trackways |
| <i>Poa pratensis</i> | Smooth Meadow-grass | Common. Meadows |
| <i>Poa nemoralis</i> | Wood Meadow-grass | Frequent |
| <i>Poa supina</i> | Creeping Meadow-grass | Frequent |
| <i>Poa trivialis</i> | Rough Meadow-grass | Frequent |
| <i>Schedonorus pratensis</i> | Meadow Fescue | Common |
| <i>Sesleria caerulea</i> | Blue Moor-grass | Common. Limestone pasture |
| Tofieldiaceae | Scottish Asphodel Family | |
| <i>Tofieldia calyculata</i> | Tofield's Asphodel | Common |

Birds (✓=recorded but not counted; H = heard only)

| | Common name | Latin name | June/July | | | | | | | |
|---|---------------------|---------------------------|-----------|----|----|----|----|----|----|---|
| | | | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 1 |
| 1 | Mute Swan | <i>Cygnus olor</i> | ✓ | | | | | | | |
| 2 | Great Crested Grebe | <i>Podiceps cristatus</i> | ✓ | | | | | | | |
| 3 | Red Kite | <i>Milvus milvus</i> | ✓ | | | | | ✓ | ✓ | |
| 4 | Black Kite | <i>Milvus migrans</i> | ✓ | | | | | | | |
| 5 | Bearded Vulture | <i>Gypaetus barbatus</i> | | | ✓ | | | | | |
| 6 | Common Buzzard | <i>Buteo buteo</i> | ✓ | | | | ✓ | ✓ | ✓ | |

| | Common name | Latin name | June/July | | | | | | | |
|----|---------------------------|-----------------------------------|-----------|----|----|----|----|----|----|---|
| | | | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 1 |
| 7 | Eurasian Coot | <i>Fulica atra</i> | ✓ | | | | | | | |
| 8 | Black-headed Gull | <i>Chroicocephalus ridibundus</i> | ✓ | | | | | | | |
| 9 | Yellow-legged Gull | <i>Larus michahellis</i> | ✓ | | | | | | | |
| 10 | Common Wood Pigeon | <i>Columba palumbus</i> | ✓ | | | | | | | |
| 11 | Eurasian Collared Dove | <i>Streptopelia decaocto</i> | ✓ | | | | | | ✓ | |
| 12 | Common Swift | <i>Apus apus</i> | ✓ | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 13 | Black Woodpecker | <i>Dryocopus martius</i> | | | | ✓ | | | | |
| 14 | Eurasian Jay | <i>Garrulus glandarius</i> | | | | ✓ | ✓ | | | |
| 15 | Eurasian Magpie | <i>Pica pica</i> | | | | | | | ✓ | |
| 16 | Spotted Nutcracker | <i>Nucifraga caryocatactes</i> | | | ✓ | ✓ | ✓ | | | |
| 17 | Alpine Chough | <i>Pyrrhocorax graculus</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 18 | Carrion Crow | <i>Corvus corone</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 19 | Northern Raven | <i>Corvus corax</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 20 | European Crested Tit | <i>Lophophanes cristatus</i> | | | | ✓ | | | | |
| 21 | Great Tit | <i>Parus major</i> | | | | ✓ | ✓ | ✓ | ✓ | |
| 22 | Barn Swallow | <i>Hirundo rustica</i> | ✓ | | | | | | ✓ | |
| 23 | Common House Martin | <i>Delichon urbicum</i> | ✓ | | | ✓ | ✓ | ✓ | | |
| 24 | Willow Warbler | <i>Phylloscopus trochilus</i> | | | | | H | | | |
| 25 | Common Chiffchaff | <i>Phylloscopus collybita</i> | | | | | H | | H | |
| 26 | Eurasian Blackcap | <i>Sylvia atricapilla</i> | | | | H | H | | | |
| 27 | Goldcrest | <i>Regulus regulus</i> | | | | H | | | | |
| 28 | Eurasian Wren | <i>Troglodytes troglodytes</i> | | | | H | | | | |
| 29 | Eurasian Treecreeper | <i>Certhia familiaris</i> | | | | ✓ | | | | |
| 30 | Ring Ouzel | <i>Turdus torquatus</i> | | ✓ | | | | | | |
| 31 | Common Blackbird | <i>Turdus merula</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 32 | Fieldfare | <i>Turdus pilaris</i> | | | | | ✓ | | | |
| 33 | Mistle Thrush | <i>Turdus viscivorus</i> | | | | ✓ | | | | |
| 34 | Rufous-tailed Rock Thrush | <i>Monticola saxatilis</i> | | | | | ✓ | | | |
| 35 | European Robin | <i>Erithacus rubecula</i> | H | | | | | | | |
| 36 | Black Redstart | <i>Phoenicurus ochruros</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 37 | Northern Wheatear | <i>Oenanthe oenanthe</i> | | ✓ | | | ✓ | | ✓ | |
| 38 | House Sparrow | <i>Passer domesticus</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| 39 | White-winged Snowfinch | <i>Montifringilla nivalis</i> | | ✓ | | | | | | |
| 40 | Alpine Accentor | <i>Prunella collaris</i> | | ✓ | | | | | | ✓ |
| 41 | Dunnock | <i>Prunella modularis</i> | | | | | H | | | |
| 42 | White Wagtail | <i>Motacilla alba</i> | ✓ | ✓ | | ✓ | ✓ | ✓ | | |
| 43 | Water Pipit | <i>Anthus spinoletta</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 44 | Common Chaffinch | <i>Fringilla coelebs</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 45 | European Serin | <i>Serinus serinus</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | | |
| 46 | European Greenfinch | <i>Chloris chloris</i> | H | H | | H | | | | |
| 47 | European Goldfinch | <i>Carduelis carduelis</i> | | | | ✓ | | ✓ | | |
| 48 | Common Linnet | <i>Linaria cannabina</i> | | ✓ | ✓ | | | | | |
| 49 | Eurasian Bullfinch | <i>Pyrrhula pyrrhula</i> | | | | ✓ | H | | | |

Mammals (E=evidence of)

| | | | | | | | | | | |
|---|----------------|----------------------------|--|---|---|---|---|---|---|--|
| 1 | Mole | <i>Talpa europaea</i> | | E | | | | E | | |
| 2 | Red Squirrel | <i>Sciurus vulgaris</i> | | | | ✓ | | ✓ | | |
| 3 | Alpine Marmot | <i>Marmota marmota</i> | | ✓ | | | ✓ | | H | |
| 4 | Snow Vole | <i>Microtus nivalis</i> | | E | | | | | | |
| 5 | Alpine Ibex | <i>Capra ibex</i> | | | ✓ | | | | | |
| 6 | Alpine Chamois | <i>Rupicapra rupicapra</i> | | ✓ | | ✓ | | | | |

| | Common name | Latin name | June/July | | | | | | |
|--|-------------|------------|-----------|----|----|----|----|----|----|
| | | | 24 | 25 | 26 | 27 | 28 | 29 | 30 |

Amphibians (D = Dead)

| | | | | | | | | | | |
|---|-------------------|--------------------------|--|---|---|--|---|--|--|--|
| 1 | Common Frog | <i>Rana temporaria</i> | | | ✓ | | | | | |
| 2 | Common Newt | <i>Triturus vulgaris</i> | | ✓ | ✓ | | | | | |
| 3 | Alpine Salamander | <i>Salamandra atra</i> | | D | | | D | | | |

Butterflies

| | | | | | | | | | | |
|----|---------------------------------|-------------------------------|--|---|---|---|---|---|---|--|
| 1 | Swallowtail | <i>Papilio machaon</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 2 | Black-veined White | <i>Aporia crataegi</i> | | | | ✓ | ✓ | | ✓ | |
| 3 | Large White | <i>Pieris brassicae</i> | | | | | | ✓ | | |
| 4 | Small White | <i>Pieris rapae</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 5 | Mountain Green-veined White | <i>Pieris bryoniae</i> | | | ✓ | ✓ | | | | |
| 6 | Peak White | <i>Pontia callidice</i> | | | | | | | ✓ | |
| 7 | Orange Tip | <i>Anthocharis cardamines</i> | | | ✓ | | ✓ | | | |
| 8 | Clouded Yellow | <i>Colias crocea</i> | | | | | | ✓ | | |
| 9 | Mountain Clouded Yellow | <i>Colias phicomone</i> | | | ✓ | ✓ | | | | |
| 10 | Wood White | <i>Leptidea sinapis</i> | | | ✓ | | | | | |
| 11 | Green Hairstreak | <i>Callophrys rubi</i> | | | ✓ | | | ✓ | | |
| 12 | Little Blue | <i>Cupido minimus</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 13 | Alpine Argus | <i>Albulina orbitulus</i> | | | | | | | ✓ | |
| 14 | Common Blue | <i>Polyommatus icarus</i> | | ✓ | ✓ | ✓ | ✓ | | ✓ | |
| 15 | Painted Lady | <i>Cynthia cardui</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 16 | Small Tortoiseshell | <i>Aglais urticae</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 17 | Small Pearl-bordered Fritillary | <i>Clossiana selene</i> | | | | | ✓ | ✓ | ✓ | |
| 18 | Pearl-bordered Fritillary | <i>Clossiana euphrosyne</i> | | | | | ✓ | ✓ | ✓ | |
| 19 | Marbled White | <i>Melanargia galathea</i> | | | | ✓ | | | | |
| 20 | Meadow Brown | <i>Maniola jurtina</i> | | | ✓ | ✓ | ✓ | | | |
| 21 | Alpine Heath | <i>Coenonympha gardetta</i> | | | ✓ | ✓ | | | ✓ | |
| 22 | Speckled Wood | <i>Pararge aegeria</i> | | | | ✓ | ✓ | | | |
| 23 | Dingy Skipper | <i>Erynnis tages</i> | | | ✓ | | | | | |
| 24 | Essex Skipper | <i>Thymelicus lineolus</i> | | | | ✓ | ✓ | | | |
| 25 | Grizzled Skipper | <i>Pyrgus malvae</i> | | | ✓ | | | | | |

Moths (L=larva)

| | | | | | | | | | | |
|---|----------------------|---------------------------------|--|---|---|---|---|---|---|--|
| 1 | Hummingbird Hawkmoth | <i>Macroglossum stellatarum</i> | | ✓ | | ✓ | ✓ | ✓ | ✓ | |
| 2 | Chimney Sweeper | <i>Odezia atrata</i> | | | | | ✓ | | | |
| 3 | Lackey Moth | <i>Malacosoma neustria</i> | | | L | | | | | |
| 4 | Latticed Heath | <i>Chiasmia clathrata</i> | | | | | | ✓ | | |

Dragonflies

| | | | | | | | | | | |
|---|--------------------|---------------------------|--|--|---|--|--|--|--|--|
| 1 | White-faced Darter | <i>Leucorrhinia dubia</i> | | | ✓ | | | | | |
|---|--------------------|---------------------------|--|--|---|--|--|--|--|--|

Others

| | | | | | | | | | | |
|---|-----------------------|------------------------|--|---|--|--|---|--|---|--|
| 1 | An alpine leaf beetle | <i>Oreina gloriosa</i> | | ✓ | | | ✓ | | ✓ | |
|---|-----------------------|------------------------|--|---|--|--|---|--|---|--|