

Wengen – Alpine Flowers of the Swiss Alps

Naturetrek Tour Report

25th June – 2nd July 2025



Cerastium latifolium



Rhododendron ferrugineum



Noccaea rotundifolia



Oxytropis campestris

Tour report and photos by David Tattersfield



Naturetrek Mingledown Barn Wolf's Lane Chawton Alton Hampshire GU34 3HJ UK

T: +44 (0)1962 733051

E: info@naturetrek.co.uk

W: www.naturetrek.co.uk

Tour participants: David Tattersfield (tour leader and botanist) with 13 Naturetrek clients.

Day 1

Wednesday 25th June

After assembling at Zurich airport, we boarded the train to Interlaken. However, the journey did not go quite to plan. All trains, beyond Zurich main station, had been cancelled owing to an interruption in the overhead power supply. After some confusion and a short wait, we transferred to another train, which took us on a much longer route via Lucerne and Brienz. At Interlaken, we quickly transferred to the regional train to Lauterbrunnen, and eventually arrived in Wengen at 21.45. After a very long day, we appreciated the cold buffet, that awaited us, before retiring to bed for much-needed sleep.

Day 2

Thursday 26th June

On a sunny morning, we set off to walk down to Lauterbrunnen. The forest understorey harboured plants more typical of lower elevations, such as Dog's Mercury *Mercurialis perennis*, Nettle-leaved Speedwell *Veronica urticifolia*, Cowslip *Primula veris*, Herb-Paris *Paris quadrifolia*, and Whorled Solomon's-seal *Polygonatum verticillatum*. In open glades, we added Nettle-leaved Bellflower *Campanula trachelium*, Dark Columbine *Aquilegia atrata*, the large Cabbage Thistle *Cirsium oleraceum*, and the tall, white plumes of Goat's-beard Spiraea *Aruncus dioicus*. Following coffee in Lauterbrunnen, we caught the bus to Trummelbach and enjoyed a picnic lunch, beside the river.

A thunderstorm arrived, just as we had finished eating, and we crossed the road to the shelter of the Trummelbach Falls. Using the lift and tunnels, we visited all 10 of the accessible waterfalls, a truly amazing experience. As we emerged, the rain was easing, and we walked down past the lower falls, finding Branched St Bernard's Lily *Anthericum ramosum*, Mountain Germander *Teucrium montanum*, and the rhizomatous Mountain Bladder-fern *Cystopteris montana*.

After drinks at the café, we enjoyed a leisurely walk down the valley, past the impressive Staubach Falls, to Lauterbrunnen, where we caught the 17.30 train, back to Wengen. After dinner, some of us enjoyed a folk evening, provided by the local community.

Day 3

Friday 27th June

After leaving the train at Wengernalp, we followed the road down to Wixi, through colourful meadows, full of Greater Yellow-rattle *Rhinanthus alectorolophus*, Common Bistort *Bistorta officinalis*, Spiked Rampion *Phyteuma spicatum*, and Wood Crane's-bill *Geranium sylvaticum*. In wetter places, the dark purple Broad-leaved Marsh-orchid *Dactylorhiza majalis* was common, along with Twayblade *Neottia ovata*, and deep-blue Bavarian Gentian *Gentiana bavarica*. Around a corner, we found colourful plants on the limestone rocks, such as Cypress Spurge *Euphorbia cyparissias*, Horseshoe Vetch *Hippocrepis comosa*, the pale blue Matted Globularia *Globularia cordifolia* and the grey-leaved Chamois Ragwort *Senecio doronicum*, while in shady areas of the forest we found Serrated Wintergreen *Orthilia secunda*, One-flowered Wintergreen *Moneses uniflora* and the parasitic Coralroot Orchid *Corallorhiza trifida*.

At Biglenalp, we relaxed, over lunch, at a lovely viewpoint, beside a stream, before exploring the surrounding pasture. A diversity of new species included Alpine Calamint *Clinopodium alpinum*, the soft-pink Mountain Sainfoin

Onobrychis montana, large blue flowers of Alpine Columbine *Aquilegia alpina*, and a single well-flowered plant of Lady's-slipper *Cypripedium calceolus*.

During the afternoon, we gradually descended through broad meadows, where the tall yellow sentinels of Great Yellow Gentian *Gentiana lutea* lit up the scene. Patches of damp soil were attracting Mazarine Blue, Chequered Skipper and the attractive day-flying moth, White-spotted Sable, and around a small pond we found Broad-bodied and Four-spotted Chasers. On the edge of the forest, a close encounter with a family of Northern Nutcrackers, was another highlight and on the damp floor of the forest, we found colonies of the tiny Lesser Twayblade *Neottia cordata*, almost invisible against the mossy background. A broad farm track led us past the cowsheds at Mettlenalp and down through the forest, where damp ditches contained Round-leaved Saxifrage *Saxifraga rotundifolia*, Yellow Mountain Saxifrage *Saxifraga aizoides*, Starry Saxifrage *Micranthes stellaris* and Common Butterwort *Pinguicula vulgaris*. After a hot day and a longish walk, it was a relief to catch the train from Almend, back to Wengen.

Day 4

Saturday 28th June

After a short cable-car ride to Mannlichen, we were surrounded by stunning scenery, with the famous trio of peaks, the Jungfrau, Monch and Eiger, dominating the skyline, to the south. We enjoyed a busy morning of botanical discoveries, as we made our way to the summit. Particularly eye-catching were Long-spurred Pansy *Viola calcarata*, Bird's-eye Primrose *Primula farinosa*, Trumpet Gentian *Gentiana acaulis*, Spring Gentian *Gentiana verna*, and the pale yellow Alpine Pasqueflower *Pulsatilla alpina* subsp. *apiifolia*. Where snow had recently melted, we found yellow Alpine Avens *Geum montanum*, Mountain Buttercup *Ranunculus montanus*, the white Kupfer's Buttercup *Ranunculus kuepferi*, and Alpine Snowbell *Soldanella alpina*. The season was well advanced and Small-white Orchid *Pseudorchis albida*, Alpine Vanilla Orchid *Nigritella rhellicani*, and Fragrant Orchid *Gymnadenia conopsea*, were already coming into flower. Just beyond the summit, on a north-facing slope, we found Crimson-tipped Lousewort *Pedicularis oederi*, growing among dwarf shrubs of Mountain Avens *Dryas octopetala*, Retuse-leaved Willow *Salix retusa*, Net-leaved Willow *Salix reticulata* and Cowberry *Vaccinium vitis-idaea*.

Returning down the mountain, we relaxed over coffee at the Mannlichen Berghaus, before having our lunch on a grassy knoll, overlooking Grindewald. White-winged Snowfinch foraged for seed, nearby, and around us a new range of plants, growing on the damp acidic shales, included Shaggy Hawkweed *Hieracium villosum*, Cat's-foot *Antennaria dioica*, the scented Musk Milfoil *Achillea erba-rotta* subsp. *moschata*, Mossy Saxifrage *Saxifraga bryoides*, and large bushes of Alpenrose *Rhododendron ferrugineum*.

During the remainder of the afternoon, we followed an easy trail down to Kleine Scheidegg. Lots of flowers lined our route, with notable highlights of Large-flowered Leopard's-bane *Doronicum grandiflorum*, Oxlip *Primula elatior*, and Pallid Milk-vetch *Astragalus frigidus*. The steep north face of the Eiger grew ever closer, across the valley, and around snow-beds on the Lauberhorn, we picked out no less than 24 Alpine Chamois. After finishing our walk, we caught the next train to Wengen. After dinner on the terrace, a pink alpenglow on the Jungfrau brought the day to a close.

Day 5

Sunday 29th June

Taking an early train to Lauterbrunnen, we rode the cable-car to Grutschalp and boarded another train to Murren. After a short funicular ride, we enjoyed a coffee break at Almendhubel, before setting off to explore. The vegetation

on this side of the valley provided a complete contrast to that of previous days. Tall herbs, such as White False-helleborine *Veratrum album*, *Adenostyles alliariae*, Masterwort *Peucedanum ostruthium*, and the reddish-purple *Gentiana purpurea* were common, among thickets of Green Alder *Alnus alnobetula* subsp. *alnobetula*, and a luxuriant growth of ferns. The bright-yellow Mountain Pansy *Viola lutea* was plentiful, and in marshy ground, a diverse array of orchids included Fragrant Orchid, Twayblade, robust plants of Early-purple Orchid *Orchis mascula* and Round-headed Orchid *Traunsteinera globosa*.

During lunch, we enjoyed superb views of the panorama of peaks across the valley. Afterwards, some of the group walked down an easy track to Winteregg, while rest of us followed the aptly named Mountain View Trail. We crossed heather-clad moorland, where White-faced Darters were flying around acidic pools, among the fluffy white flowers of Scheuchzer's Cotton-grass *Eriophorum scheuchzeri*. During our descent, notable finds included the pale-yellow Alpine Leek *Allium victorialis*, a single plant of the rare Alpine Bladder-fern *Cystopteris alpina*, large stands of Alpine Blue-sowthistle *Cicerbita alpina*, and the delicate Lesser Astartia *Astartia minor*. A final steep path, through the forest, returned us to Grutschalp, and our return transport to Wengen.

Day 6

Monday 30th June

After travelling down to Wilderswil, we boarded the delightful old train for the steep climb to Schynige Platte. After coffee, we spent the morning exploring the many delights of the Alpine Garden, where the natural setting has been thoughtfully augmented by the addition of new habitats and plants from other regions of the Swiss Alps. As usual, the garden reflected the skills and dedication of its staff and was looking at its very best.

After lunch on the station platform, some of our number returned to the garden, while the rest of us followed a panoramic walk around the limestone pinnacles of Gaube and the Oberberghorn. When we reached the ridge, the weather took a turn for the worse, and we watched curtains of heavy rain falling over the Brienzensee, far below us. The atmospherics, however, did not distract us, and we enjoyed a wonderful selection of colourful plants. Among them were large patches of Great Yellow Gentian, the striking Yellow Pea *Lathyrus laevigatus* subsp. *occidentalis*, Yellow Bellflower *Campanula thyrsoides*, and Starry Hare's-ear *Bupleurum stellatum*. On the cliffs, we found deep-purple Mountain Milk-vetch *Oxytropis jacquinii*, Alpine Aster *Aster alpinus*, Wood Pink *Dianthus sylvestris*, and Dwarf Buckthorn *Rhamnus pumilus*. A striking feature of the Oberberghorn is the limestone scree, that lies at a steep angle below the cliffs, colonised by extensive patches of Limestone Fern *Gymnocarpium robertianum*, Angular Solomon's-seal *Polygonatum odoratum*, Whorled Solomon's-seal *Polygonatum verticillatum*, and Lily-of-the-Valley *Convallaria majalis*.

As the rain became heavier, we made a dash for the station and returned to Wilderswil, where we enjoyed an ice-cream, while waiting for our connecting train.

Day 7

Tuesday 1 July

On our final day, we took the trains to Kleine Scheidegg and on to the Eigergletscher station. Above the station platform, we made our way onto the moraines, below the west ridge of the Eiger. The jumble of frost-shattered rocks appeared barren, at first glance, but we soon found some of the colourful, pioneer plants that colonise this harsh environment. Most attractive, among these, were Fairy's-thimble *Campanula cochleariifolia*, Mt Cenis Pansy *Viola cenisia*, Broad-leaved Mouse-ear *Cerastium latifolium*, Chamois Cress *Hornungia alpina* subsp. *alpina*, and the

sweetly scented Round-leaved Penny-cress *Noccaea rotundifolia*. We followed the narrow ridge of the medial moraine back towards the station, where the slow process of colonisation was more advanced and the ground was covered with sprawling mats of Retuse-leaved Willow and large cushions of Moss Campion *Silene acaulis*, interspersed with Spring Gentians, and a host of other plants.

Alpine Chough provided us with great entertainment, while we enjoyed coffee and cakes, on the terrace of the station café. Around a corner, at the start of the Eiger Trail, Large-flowered Leopard's-bane, Short-leaved Gentian *Gentiana brachyphylla*, and Glacier Crowfoot *Ranunculus glacialis* were among several plants, starting to colonise the steep north-facing screes. Below the station, a grassy slope provided an ideal spot for our lunch, and once again, we attracted the attention of dozens of Alpine Chough.

Most of the group walked down to Kleine Scheidegg to catch the train, leaving four of us to continue down the moraine and follow a steep trail to Wixi, and, finally, Wengernalp.

Day 8

Wednesday 2nd July

Sadly, our wonderful week had come to an end. After breakfast, we thanked Martina and her staff for their excellent hospitality, before walking down to the station for our return journey to Zurich.

Receive our e-newsletter

Join the Naturetrek email list and be the first to hear about new tours, additional departures and new dates, tour reports and special offers.

naturetrek.co.uk/e-newsletter-signup

Scan to
sign up!



Social Media

We're social! Follow us on Instagram, Facebook, Bluesky and X and be the first to hear about the launch of new tours, offers, exciting sightings and photos from our recently returned holidays.



[instagram.com/naturetrek_wildlife_holidays](https://www.instagram.com/naturetrek_wildlife_holidays)



[facebook.com/naturetrekwildlifeholidays](https://www.facebook.com/naturetrekwildlifeholidays)



[naturetrek.bsky.social](https://www.bsky.social/naturetrek)



x.com/naturetrektours (formerly Twitter)

Species lists

Plants

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|----------------------------------|------------------------------|-------------------------------|
| PTERIDOPHYTES | FERNS AND FERN ALLIES | |
| Aspleniaceae | Spleenwort Family | |
| <i>Asplenium adiantum-nigrum</i> | Black Spleenwort | Shady cliffs |
| <i>Asplenium ruta-muraria</i> | Wall-rue | Walls |
| <i>Asplenium scolopendrium</i> | Hart's-tongue | Lowland woods |
| <i>Asplenium septentrionale</i> | Forked Spleenwort | Rare, on rocks |
| <i>Asplenium trichomanes</i> | Maidenhair Spleenwort | Walls and cliffs |
| <i>Asplenium viride</i> | Green Spleenwort | Mountain rocks |
| Athyriaceae | Lady-fern Family | |
| <i>Athyrium distentifolium</i> | Alpine Lady-fern | Mountain woods. Above Murren |
| <i>Athyrium filix-femina</i> | Lady-fern | Occasional |
| Blechnaceae | Hard-fern Family | |
| <i>Blechnum spicant</i> | Hard-fern | Locally common |
| Cystopteridaceae | Bladder-fern Family | |
| <i>Cystopteris alpina</i> | Alpine Bladder-fern | Rare. Grutschalp |
| <i>Cystopteris fragilis</i> | Brittle Bladder-fern | Frequent on rocks |
| <i>Cystopteris montana</i> | Mountain Bladder-fern | Local. Trummelbach |
| <i>Gymnocarpium dryopteris</i> | Oak Fern | Woods and rocks. Local |
| <i>Gymnocarpium robertianum</i> | Limestone Fern | Limestone rocks. Oberberghorn |
| Dennstaedtiaceae | Bracken Family | |
| <i>Pteridium aquilinum</i> | Bracken | Valley woodland |
| Dryopteridaceae | Buckler-fern Family | |
| <i>Dryopteris affinis</i> | Scaly Male-fern | Widespread |
| <i>Dryopteris carthusiana</i> | Narrow Buckler-fern | Damp scrub |
| <i>Dryopteris dilatata</i> | Broad Buckler-fern | Frequent |
| <i>Dryopteris expansa</i> | Northern Buckler-fern | Damp scrub |
| <i>Dryopteris filix-mas</i> | Male-fern | Widespread |
| <i>Dryopteris submontana</i> | Rigid Buckler-fern | Schynige Platte |
| <i>Polystichum aculeatum</i> | Hard Shield-fern | Woods. Occasional |
| <i>Polystichum lonchitis</i> | Holly-fern | Mountain rocks |
| Equisetaceae | Horsetail Family | |
| <i>Equisetum arvense</i> | Field Horsetail | Meadows and tracksides |
| <i>Equisetum palustre</i> | Marsh Horsetail | Wet pasture |
| <i>Equisetum sylvaticum</i> | Wood Horsetail | Woods and damp pasture |
| Lycopodiaceae | Clubmoss Family | |
| <i>Huperzia selago</i> | Fir Clubmoss | Frequent. Acidic pasture |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|-------------------------------|------------------------------|
| <i>Lycopodium alpinum</i> | Alpine Clubmoss | Occasional. Mountain turf |
| <i>Lycopodium annotinum</i> | Interrupted Clubmoss | Local. Almend |
| <i>Lycopodium clavatum</i> | Stag's-horn Clubmoss | Local. Above Grutschalp |
| Ophioglossaceae | Adder's-tongue Family | |
| <i>Botrychium lunaria</i> | Moonwort | Short turf. Widespread |
| Polypodiaceae | Polypody Family | |
| <i>Polypodium vulgare</i> | Common Polypody | Occasional on rocks |
| Pteridaceae | Maidenhair Fern Family | |
| <i>Cryptogramma crispera</i> | Parsley Fern | Acidic rocks |
| Selaginellaceae | Lesser Clubmoss Family | |
| <i>Selaginella selaginoides</i> | Lesser Clubmoss | Occasional in turf |
| Thelypteridaceae | Marsh Fern Family | |
| <i>Oreopteris limbosperma</i> | Lemon-scented Fern | Green Alder scrub and woods |
| <i>Phegopteris connectilis</i> | Beech Fern | Woods. Occasional |
| Woodsiaceae | Cliff Fern Family | |
| <i>Woodsia alpina</i> | Alpine Woodsia | Rare. Almendhubel |
| GYMNOSPERMS | CONIFERS | |
| Cupressaceae | Juniper Family | |
| <i>Juniperus communis</i> var. <i>saxatilis</i> | Common Juniper | Mountain pasture |
| Pinaceae | Pine Family | |
| <i>Abies alba</i> | European Silver-fir | Occasional. Forests |
| <i>Larix decidua</i> | Larch | Occasional |
| <i>Picea abies</i> | Norway Spruce | Dominant. Forests |
| <i>Pinus cembra</i> | Arolla Pine | Locally dominant at treeline |
| <i>Pinus sylvestris</i> | Scots Pine | Occasional |
| <i>Pseudotsuga menziesii</i> ‡ | Douglas Fir | Forest below Wengen |
| ANGIOSPERMS | FLOWERING PLANTS | |
| DICOTYLEDONS | | |
| Amaranthaceae | Goosefoot Family | |
| <i>Blitum bonus-henricus</i> | Good-King-Henry | Frequent weed |
| Apiaceae | Carrot Family | |
| <i>Aegopodium podagraria</i> | Ground Elder | Below Wengen |
| <i>Angelica sylvestris</i> | Wild Angelica | Common in lowlands |
| <i>Anthriscus sylvestris</i> | Cow Parsley | Frequent |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|---------------------------|------------------------------|
| <i>Astrantia major</i> | Astrantia | Meadows |
| <i>Astrantia minor</i> | Lesser Astrantia | Locally common |
| <i>Athamanta cretensis</i> | Athamanta | Rocky places and moraines |
| <i>Bupleurum stellatum</i> | Starry Hare's-ear | Mountain rocks |
| <i>Carum carvi</i> | Caraway | Tracksides |
| <i>Chaerophyllum hirsutum</i> | Hairy Chervil | Woods |
| <i>Daucus carota</i> | Wild Carrot | Wilderswil |
| <i>Heracleum sphondylium</i> | Hogweed | Common |
| <i>Laserpitium halleri</i> | Haller's Sermountain | Occasional |
| <i>Laserpitium latifolium</i> | Broad-leaved Sermountain | Trummelbach |
| <i>Mutellina adonidifolia</i> | Alpine Lovage | Mountain pasture |
| <i>Neogaya simplex</i> | Unbranched Lovage | High-mountain pasture |
| <i>Peucedanum ostruthium</i> | Masterwort | Common |
| <i>Pimpinella major</i> | Greater Burnet-saxifrage | Common |
| <i>Pimpinella saxifraga</i> | Burnet-saxifrage | Meadows |
| <i>Sanicula europaea</i> | Sanicle | Lowland woods |
| <i>Siler montanum</i> | Narrow-leaved Sermountain | Occasional |
| Apocynaceae | | |
| <i>Vincetoxicum hirundinaria</i> | Swallow-wort | Frequent |
| Araliaceae | | |
| <i>Hedera helix</i> | Ivy | Occasional |
| Asteraceae | | |
| <i>Adenostyles alliariae</i> | Adenostyles | Green Alder scrub |
| <i>Adenostyles alpina</i> subsp. <i>alpina</i> | Hairless Adenostyles | Screes. Oberberghorn |
| <i>Adenostyles leucophylla</i> | White-leaved Adenostyles | Screes. Oberberghorn |
| <i>Achillea atrata</i> | Dark-stemmed Sneezewort | Frequent on limestone |
| <i>Achillea erba-rotta</i> subsp. <i>moschata</i> | Musk Milfoil | Acidic mountain pasture |
| <i>Achillea macrophylla</i> | Large-leaved Yarrow | Almendhubel |
| <i>Achillea millefolium</i> | Yarrow | Meadows |
| <i>Antennaria carpatica</i> | Carpathian Cat's-foot | Occasional. Mountain pasture |
| <i>Antennaria dioica</i> | Cat's-foot | Common |
| <i>Arnica montana</i> | Arnica | Common |
| <i>Aster alpinus</i> | Alpine Aster | Mountain rocks and turf |
| <i>Aster bellidiastrum</i> | False Aster | Common |
| <i>Bellis perennis</i> | Daisy | Frequent |
| <i>Carduus defloratus</i> | Alpine Thistle | Common |
| <i>Carduus personata</i> | Great Marsh Thistle | Murren |
| <i>Carlina acaulis</i> | Stemless Carlina Thistle | Mountain pasture |
| <i>Centaurea montana</i> | Mountain Cornflower | Meadows |
| <i>Centaurea scabiosa</i> | Greater Knapweed | Meadows |
| <i>Cicerbita alpina</i> | Alpine Blue-sowthistle | Forests and scrub |
| <i>Cirsium acaule</i> | Dwarf Thistle | Mountain pasture |
| <i>Cirsium oleraceum</i> | Cabbage Thistle | Common below Wengen |
| <i>Cirsium palustre</i> | Marsh Thistle | Occasional. Wet places |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---------------------------------|-------------------------------|---------------------------|
| <i>Cirsium spinosissimum</i> | Spiniest Thistle | Mountain pasture |
| <i>Crepis aurea</i> | Golden Hawk's-beard | Meadows |
| <i>Crepis paludosa</i> | Marsh Hawk's-beard | Wet meadows |
| <i>Crepis pontana</i> | Mountain Hawk's-beard | Mountain pasture |
| <i>Crepis pyrenaica</i> | Pyrenean Hawk's-beard | Meadows. Murren |
| <i>Doronicum grandiflorum</i> | Large-flowered Leopard's-bane | Scree |
| <i>Erigeron alpinus</i> | Alpine Fleabane | Mountain rocks. Local |
| <i>Erigeron annuus</i> ‡ | Tall Fleabane | Wilderswil |
| <i>Eupatorium cannabinum</i> | Hemp Agrimony | Lowland |
| <i>Hieracium angustifolium</i> | Glacier Hawkweed | Occasional |
| <i>Hieracium murorum</i> | Few-leaved Hawkweed | Occasional |
| <i>Hieracium piliferum</i> | Hairy Hawkweed | Occasional |
| <i>Hieracium pilosum</i> | Fimbriate-pitted Hawkweed | Occasional |
| <i>Hieracium villosum</i> | Shaggy Hawkweed | Mannlichen |
| <i>Homogyne alpina</i> | Alpine Colt's-foot | Mountain turf. Common |
| <i>Hypochaeris uniflora</i> | Giant Cat's-ear | Mannlichen |
| <i>Lapsana communis</i> | Nipplewort | Below Wengen |
| <i>Leontodon hispidus</i> | Alpine Hawkbit | Acidic pasture |
| <i>Leucanthemopsis alpina</i> | Alpine Moon Daisy | Acidic mountain turf |
| <i>Leucanthemum adustum</i> | Mountain Ox-eye | Mountain pasture |
| <i>Leucanthemum vulgare</i> | Ox-eye Daisy | Meadows |
| <i>Matricaria discoidea</i> ‡ | Pineappleweed | Ruderal |
| <i>Mycelis muralis</i> | Wall Lettuce | Walls |
| <i>Omalotheca norvegica</i> | Norwegian Cudweed | Occasional |
| <i>Omalotheca supina</i> | Dwarf Cudweed | Snowbeds |
| <i>Petasites albus</i> | White Butterbur | Mannlichen |
| <i>Petasites paradoxus</i> | Alpine Butterbur | Mountain pasture |
| <i>Pilosella aurantiaca</i> ‡ | Fox-and-cubs | Wengen |
| <i>Pilosella lactucella</i> | Glaucous Fox-and-cubs | Common |
| <i>Pilosella officinarum</i> | Mouse-ear Hawkweed | Common |
| <i>Prenanthes purpurea</i> | Purple Lettuce | Common. Forests |
| <i>Saussurea alpina</i> | Alpine Saw-wort | Mountain turf |
| <i>Scorzoneroides helvetica</i> | Swiss Hawkbit | Mountain turf. Common |
| <i>Senecio doronicum</i> | Chamois Ragwort | Limestone rocks and scree |
| <i>Senecio ovatus</i> | Wood Ragwort | Forests |
| <i>Solidago virgaurea</i> | Goldenrod | Mountain pasture |
| <i>Sonchus asper</i> | Prickly Sow-thistle | Ruderal |
| <i>Sonchus oleraceus</i> | Smooth Sow-thistle | Ruderal |
| <i>Taraxacum alpinum</i> | Alpine Dandelion | Scree |
| <i>Taraxacum officinale</i> | Dandelion | Widespread |
| <i>Tolpis staticifolia</i> | Statice-leaved Hawkweed | Alluvium and scree |
| <i>Tragopogon pratensis</i> | Goat's-beard | Meadows |
| <i>Tussilago farfara</i> | Colt's-foot | Damp places. Widespread |
| Balsaminaceae | Balsam Family | |
| <i>Impatiens parviflora</i> ‡ | Small Balsam | Woods below Wengen |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|--|--------------------------------|---------------------------|
| Betulaceae | | |
| Birch Family | | |
| <i>Alnus alnobetula</i> subsp. <i>alnobetula</i> | Green Alder | Wet mountain slopes |
| <i>Alnus glutinosa</i> | Common Alder | Valleys, by rivers |
| <i>Alnus incana</i> | Grey Alder | Valleys, by rivers |
| <i>Betula pendula</i> | Silver Birch | Widespread |
| <i>Betula pubescens</i> | Downy Birch | Frequent |
| Boraginaceae | | |
| Borage Family | | |
| <i>Asperugo procumbens</i> | Madwort | Waste ground |
| <i>Echium vulgare</i> | Viper's Bugloss | Below Wengen |
| <i>Myosotis alpestris</i> | Alpine Forget-me-not | Common |
| Brassicaceae | | |
| Cabbage Family | | |
| <i>Alliaria petiolata</i> | Garlic Mustard | Tracksides |
| <i>Arabis alpina</i> subsp. <i>alpina</i> | Alpine Rock-cress | Alpine rocks |
| <i>Arabis hirsuta</i> | Hairy Rock-cress | Limestone rocks |
| <i>Biscutella laevigata</i> | Buckler Mustard | Mountain turf and moraine |
| <i>Capsella bursa-pastoris</i> | Shepherd's-purse | Widespread |
| <i>Cardamine amara</i> | Large Bitter-cress | Streamsides |
| <i>Cardamine bellidifolia</i> subsp. <i>alpina</i> | Alpine Bitter-cress | Mountain rocks |
| <i>Cardamine flexuosa</i> | Wavy Bitter-cress | Wet places |
| <i>Cardamine hirsuta</i> | Hairy Bitter-cress | Occasional |
| <i>Cardamine impatiens</i> | Narrow-leaved Bitter-cress | Damp places |
| <i>Cardamine pratensis</i> subsp. <i>pratensis</i> | Cuckooflower | Wet places |
| <i>Cardamine resedifolia</i> | Mignonette-leaved Bitter-cress | Mountain rocks |
| <i>Draba aizoides</i> | Yellow Whitlowgrass | Moraines |
| <i>Draba dubia</i> | Austrian Whitlowgrass | Rocky places |
| <i>Draba tomentosa</i> | Woolly Whitlowgrass | Rocks. Almendhubel |
| <i>Draba verna</i> | Spring Whitlowgrass | Occasional |
| <i>Homungia alpina</i> subsp. <i>alpina</i> | Chamois Cress | Mountain rocks and screes |
| <i>Kernera saxatilis</i> | Rock Kerneria | Limestone rocks. |
| <i>Noccaea perfoliata</i> | Perfoliate Penny-cress | Near habitation |
| <i>Noccaea rotundifolia</i> | Round-leaved Penny-cress | Moraines and screes |
| <i>Pseudoturritis turrita</i> | Tower Cress | Lowland rocks |
| <i>Sisymbrium austriacum</i> | Austrian Rocket | Waste places |
| Campanulaceae | | |
| Bellflower Family | | |
| <i>Campanula barbata</i> | Bearded Bellflower | Mountain turf |
| <i>Campanula cochleariifolia</i> | Fairy's-thimble | Screes |
| <i>Campanula rapunculoides</i> | Creeping Bellflower | Wilderswil |
| <i>Campanula rhomboidalis</i> | Broad-leaved Bellflower | Common in meadows |
| <i>Campanula rotundifolia</i> | Harebell | Common |
| <i>Campanula scheuchzeri</i> | Scheuchzer's Bellflower | Mountain turf |
| <i>Campanula thyrsoides</i> | Yellow Bellflower | Schynige Platte |
| <i>Campanula trachelium</i> | Nettle-leaved Bellflower | Woods |
| <i>Phyteuma betonicifolium</i> | Betony-leaved Rampion | Acidic pasture |
| <i>Phyteuma hemisphaericum</i> | Globe-headed Rampion | Mountain Rocks |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|---------------------------|----------------------------|
| <i>Phyteuma orbiculare</i> | Round-headed Rampion | Meadows. Frequent |
| <i>Phyteuma spicatum</i> | Spiked Rampion | Meadows. Frequent |
| Cannabaceae | Hop Family | |
| <i>Humulus lupulus</i> | Hop | Below Wengen |
| Caprifoliaceae | Honeysuckle Family | |
| <i>Knautia arvensis</i> | Field Scabious | Common |
| <i>Knautia dipsacifolia</i> | Wood Scabious | Common |
| <i>Lonicera alpigena</i> | Alpine Honeysuckle | Occasional. Forests |
| <i>Lonicera caerulea</i> | Blue-berried Honeysuckle | Green Alder scrub |
| <i>Lonicera nigra</i> | Black-berried Honeysuckle | Woods |
| <i>Lonicera xylostium</i> | Fly Honeysuckle | Woods |
| <i>Scabiosa columbaria</i> | Small Scabious | Common in lowlands |
| <i>Scabiosa lucida</i> | Shining Scabious | Meadows |
| <i>Succisa pratensis</i> | Devil's-bit Scabious | Damp meadows |
| <i>Valeriana dioica</i> | Marsh Valerian | Wet pasture |
| <i>Valeriana officinalis</i> | Common Valerian | Damp meadows |
| <i>Valeriana tripteris</i> | Three-leaved Valerian | Woods |
| Caryophyllaceae | Pink Family | |
| <i>Arenaria biflora</i> | Two-flowered Sandwort | Screes |
| <i>Arenaria ciliata</i> | Ciliate-leaved Sandwort | Screes |
| <i>Atocion rupestre</i> | Rock Campion | Common on rocks |
| <i>Cerastium alpinum</i> | Alpine Mouse-ear | Mountain turf |
| <i>Cerastium arvense</i> subsp. <i>strictum</i> (<i>C. elongatum</i>) | Field Mouse-ear | Frequent |
| <i>Cerastium fontanum</i> | Common Mouse-ear | Common |
| <i>Cerastium glomeratum</i> | Sticky Mouse-ear | Waste places |
| <i>Cerastium latifolium</i> | Broad-leaved Mouse-ear | Moraines and screes |
| <i>Cherleria sedoides</i> | Mossy Cyphel | Alpine turf |
| <i>Dianthus barbatus</i> ‡ | Sweet-William | Garden escape |
| <i>Dianthus carthusianorum</i> | Carthusian Pink | Almendhubel |
| <i>Dianthus superbus</i> ‡ | Large Pink | Railway banks below Almend |
| <i>Dianthus sylvestris</i> | Wood Pink | Rocks. Local |
| <i>Dichodon cerastoides</i> | Starwort Mouse-ear | Rocks and pasture |
| <i>Gypsophila repens</i> | Alpine Gypsophila | Frequent |
| <i>Moehringia ciliata</i> | Creeping Sandwort | Screes |
| <i>Moehringia muscosa</i> | Mossy Sandwort | Shady places. Local |
| <i>Sabulina verna</i> | Vernal Sandwort | Rocky places |
| <i>Sagina procumbens</i> | Procumbent Pearlwort | Common |
| <i>Saponaria ocymoides</i> | Rock Soapwort | Cliffs below Wengen |
| <i>Silene acaulis</i> | Moss Campion | Mountain turf |
| <i>Silene dioica</i> | Red Campion | Frequent. Meadows |
| <i>Silene exscapa</i> | Stemless Moss Campion | Mountain turf |
| <i>Silene flos-cuculi</i> | Ragged-Robin | Wet flushes |
| <i>Silene nutans</i> | Nottingham Catchfly | Occasional. Rocks |
| <i>Silene vulgaris</i> | Bladder Campion | Occasional. Meadows |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|--|--------------------------------|---------------------------|
| <i>Stellaria graminea</i> | Lesser Stitchwort | Occasional |
| <i>Stellaria holostea</i> | Greater Stitchwort | Occasional |
| <i>Stellaria media</i> | Common Chickweed | Occasional |
| <i>Stellaria nemorum</i> | Wood Stitchwort | Woods |
| Celastraceae | Spindle Family | |
| <i>Parnassia palustris</i> | Grass-of-Parnassus | Damp flushes |
| Cistaceae | Rockrose Family | |
| <i>Helianthemum nummularium</i> subsp. <i>grandiflorum</i> | Large-flowered Common Rockrose | Common |
| <i>Helianthemum oelandicum</i> subsp. <i>alpestre</i> | Alpine Rockrose | Frequent |
| Convolvulaceae | Bindweed Family | |
| <i>Cuscuta epithymum</i> | Common Dodder | Frequent |
| Cornaceae | Dogwood Family | |
| <i>Cornus sanguinea</i> | Dogwood | Woods |
| Corylaceae | Hazel Family | |
| <i>Corylus avellana</i> | Hazel | Woods |
| Crassulaceae | Stonecrop Family | |
| <i>Petrosedum rupestre</i> | Rock Stonecrop | Occasional. Rocks |
| <i>Phedismus spurius</i> ‡ | Caucasian Stonecrop | Wengen |
| <i>Sedum acre</i> | Biting Stonecrop | Occasional |
| <i>Sedum album</i> | White Stonecrop | Walls. Wengen |
| <i>Sedum annuum</i> | Annual Stonecrop | Mountain rocks |
| <i>Sedum alpestre</i> | Alpine Stonecrop | Mountain rocks |
| <i>Sedum atratum</i> | Dark Stonecrop | Mountain rocks |
| <i>Sedum dasyphyllum</i> | Thick-leaved Stonecrop | Walls and rocks |
| <i>Sedum hispanicum</i> ‡ | Spanish Stonecrop | Naturalised. Wengen |
| <i>Sedum rubens</i> | Reddish Stonecrop | Rocks |
| <i>Sedum sexangulare</i> ‡ | Tasteless Stonecrop | Walls. Wengen |
| <i>Sempervivum montanum</i> | Mountain Houseleek | Common on rocks |
| <i>Sempervivum tectorum</i> | Common Houseleek | Frequent on rocks |
| Ericaceae | Heather Family | |
| <i>Arctostaphylos uva-ursi</i> | Bearberry | Acidic heaths |
| <i>Arctous alpina</i> | Alpine Bearberry | Occasional. Mountain turf |
| <i>Calluna vulgaris</i> | Ling | Frequent |
| <i>Empetrum nigrum</i> subsp. <i>hermaphroditum</i> | Hermaphrodite Crowberry | Acidic heaths |
| <i>Erica carnea</i> | Spring Heath | Frequent on limestone |
| <i>Kalmia procumbens</i> | Trailing Azalea | Acidic heaths |
| <i>Moneses uniflora</i> | One-flowered Wintergreen | Forest |
| <i>Orthilia secunda</i> | Serrated Wintergreen | Forest |
| <i>Pyrola media</i> | Intermediate Wintergreen | Above Murren |
| <i>Pyrola minor</i> | Lesser Wintergreen | Forest |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|------------------------------|-----------------------------|
| <i>Rhododendron ferrugineum</i> | Alpenrose | Common |
| <i>Rhododendron hirsutum</i> | Hairy Alpenrose | Common on limestone |
| <i>Vaccinium myrtillus</i> | Bilberry | Common |
| <i>Vaccinium uliginosum</i> agg. | Bog Whortleberry | Common |
| <i>Vaccinium vitis-idaea</i> | Cowberry | Common |
| Euphorbiaceae | Spurge Family | |
| <i>Euphorbia cyparissias</i> | Cypress Spurge | Frequent on limestone |
| <i>Euphorbia helioscopia</i> | Sun Spurge | Around habitation |
| <i>Mercurialis perennis</i> | Dog's Mercury | Woods |
| Fabaceae | Pea Family | |
| <i>Anthyllis vulneraria</i> subsp. <i>alpestris</i> | Alpine Kidney Vetch | Mountain pasture |
| <i>Astragalus alpinus</i> | Alpine Milk-vetch | Mountain turf |
| <i>Astragalus australis</i> | Southern Milk-vetch | Mountain turf |
| <i>Astragalus frigidus</i> | Pallid Milk-vetch | Limestone cliffs |
| <i>Astragalus penduliflorus</i> | Mountain Lentil | Limestone cliffs |
| <i>Coronilla emerus</i> | False Senna | Occasional |
| <i>Coronilla vaginalis</i> | Small Scorpion-vetch | Cliffs below Wengen |
| <i>Hedysarum hedysaroides</i> | Alpine Sainfoin | Mountain pasture |
| <i>Hippocrepis comosa</i> | Horseshoe Vetch | Common |
| <i>Lathyrus laevigatus</i> subsp. <i>occidentalis</i> | Yellow Pea | Grutschalp, Schynige Platte |
| <i>Lathyrus latifolius</i> ‡ | Broad-leaved Everlasting-pea | Wengen |
| <i>Lathyrus pratensis</i> | Meadow Vetchling | Meadows |
| <i>Lotus alpinus</i> | Alpine Bird's-foot-trefoil | Mountain turf |
| <i>Lotus corniculatus</i> | Bird's-foot-trefoil | Meadows |
| <i>Medicago lupulina</i> | Black Medick | Common |
| <i>Medicago sativa</i> ‡ | Lucerne | Occasional weed |
| <i>Melilotus albus</i> | White Melilot | Lowland |
| <i>Melilotus indicus</i> | Small Melilot | Lowland |
| <i>Onobrychis arenaria</i> | Small Sainfoin | Occasional |
| <i>Onobrychis montana</i> | Mountain Sainfoin | Mountain pasture |
| <i>Oxytropis campestris</i> | Yellow Milk-vetch | Locally common |
| <i>Oxytropis jacquinii</i> | Mountain Milk-vetch | Moraines. Local |
| <i>Robinia pseudoacacia</i> ‡ | False Acacia | Lowlands |
| <i>Securigera varia</i> | Crown Vetch | Wilderswil |
| <i>Trifolium alpinum</i> | Alpine Clover | Common |
| <i>Trifolium badium</i> | Brown Clover | Mountain meadows |
| <i>Trifolium montanum</i> | Mountain Clover | Mountain pasture |
| <i>Trifolium pratense</i> | Red Clover | Meadows |
| <i>Trifolium repens</i> | White Clover | Frequent |
| <i>Trifolium thalii</i> | Thal's Clover | Mountain turf and scree |
| <i>Vicia cracca</i> | Tufted Vetch | Frequent |
| <i>Vicia sepium</i> | Bush Vetch | Occasional |
| <i>Vicia sylvatica</i> | Wood Vetch | Lowland woods |
| Fagaceae | Beech Family | |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|------------------------------|----------------------------|-------------------------------|
| <i>Castanea sativa</i> ‡ | Sweet Chestnut | Occasional |
| <i>Fagus sylvatica</i> | Beech | Valley forest |
| <i>Quercus petraea</i> | Sessile Oak | Frequent. Woods |
| <i>Quercus robur</i> | Pedunculate Oak | Occasional |
| Gentianaceae | | |
| Gentian Family | | |
| <i>Gentiana acaulis</i> | Trumpet Gentian | Mountain pasture |
| <i>Gentiana asclepiadea</i> | Willow Gentian | Woods on limestone |
| <i>Gentiana bavarica</i> | Bavarian Gentian | Marshes |
| <i>Gentiana brachyphylla</i> | Short-leaved Gentian | Alpine turf |
| <i>Gentiana clusii</i> | Clusius' Gentian | Limestone pasture |
| <i>Gentiana lutea</i> | Great Yellow Gentian | Mountain pasture |
| <i>Gentiana nivalis</i> | Snow Gentian | Occasional. Mountain pasture |
| <i>Gentiana purpurea</i> | Purple Gentian | Green Alder scrub and pasture |
| <i>Gentiana verna</i> | Spring Gentian | Mountain pasture |
| Geraniaceae | | |
| Crane's-bill Family | | |
| <i>Geranium pyrenaicum</i> | Pyrenean Crane's-bill | Occasional |
| <i>Geranium robertianum</i> | Herb Robert | Occasional |
| <i>Geranium sylvaticum</i> | Wood Crane's-bill | Common |
| Hypericaceae | | |
| St John's-wort Family | | |
| <i>Hypericum maculatum</i> | Imperforate St John's-wort | Frequent |
| <i>Hypericum perforatum</i> | Perforate St John's-wort | Occasional |
| Juglandaceae | | |
| Walnut Family | | |
| <i>Juglans regia</i> ‡ | Walnut | Occasional |
| Lamiaceae | | |
| Dead-nettle Family | | |
| <i>Ajuga pyramidalis</i> | Pyramidal Bugle | Locally common |
| <i>Ajuga reptans</i> | Common Bugle | Common |
| <i>Betonica officinalis</i> | Betony | Meadows |
| <i>Clinopodium alpinum</i> | Alpine Calamint | Occasional. Rocky banks |
| <i>Clinopodium vulgare</i> | Wild Basil | Frequent |
| <i>Galeopsis tetrahit</i> | Common Hemp-nettle | Occasional |
| <i>Glechoma hederacea</i> | Ground-ivy | Woods |
| <i>Lamiaeum galeobdolon</i> | Yellow Archangel | Woods |
| <i>Lamium maculatum</i> | Spotted Dead-nettle | Occasional |
| <i>Lamium purpureum</i> | Red Dead-nettle | Occasional |
| <i>Origanum vulgare</i> | Marjoram | Occasional. Meadows |
| <i>Prunella grandiflora</i> | Large-flowered Self-heal | Meadows. Trummelbach |
| <i>Prunella vulgaris</i> | Self-heal | Common |
| <i>Salvia glutinosa</i> | Sticky Clary | Forest. Wilderwil |
| <i>Salvia pratensis</i> | Meadow Clary | Meadows. Trummelbach |
| <i>Stachys alpina</i> | Limestone Woundwort | Below Wengen |
| <i>Stachys sylvatica</i> | Hedge Woundwort | Woods |
| <i>Teucrium chamaedrys</i> | Wall Germander | Below Wengen |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|--|------------------------------|------------------------------|
| <i>Teucrium montanum</i> | Mountain Germander | Trummelbach |
| <i>Teucrium scorodonia</i> | Wood Sage | Woods |
| <i>Thymus praecox</i> subsp. <i>polytrichus</i> | Wild Thyme | Common |
| Lentibulariaceae | Butterwort Family | |
| <i>Pinguicula alpina</i> | Alpine Butterwort | Mountain rocks and turf |
| <i>Pinguicula vulgaris</i> | Common Butterwort | Wet turf |
| Linaceae | Flax Family | |
| <i>Linum alpinum</i> ‡ | Alpine Flax | Wengernalp and Alpine Garden |
| <i>Linum catharticum</i> | Fairy Flax | Common |
| Malvaceae | Mallow Family | |
| <i>Tilia cordata</i> | Small-leaved Lime | Lowland |
| <i>Tilia platyphyllos</i> | Large-leaved Lime | Lowland |
| Oleaceae | Ash Family | |
| <i>Fraxinus excelsior</i> | Ash | Lowland |
| <i>Ligustrum vulgare</i> | Wild Privet | Lowland |
| Onagraceae | Willowherb Family | |
| <i>Circaea lutetiana</i> | Enchanter's-nightshade | Lauterbrunnen |
| <i>Epilobium alpestre</i> | Whorled-leaved Willowherb | Frequent |
| <i>Epilobium alsinifolium</i> | Chickweed Willowherb | Frequent |
| <i>Epilobium anagallidifolium</i> | Pimpernel-leaved Willowherb | Frequent |
| <i>Epilobium angustifolium</i> | Rosebay Willowherb | Occasional |
| <i>Epilobium hirsutum</i> | Great Willowherb | Lowlands |
| <i>Epilobium montanum</i> | Broad-leaved Willowherb | Occasional |
| Orobanchaceae | Broomrape Family | |
| <i>Bartsia alpina</i> | Alpine Bartsia | Mountain turf |
| <i>Euphrasia minima</i> | Dwarf Eyebright | Local. Mountain pasture |
| <i>Euphrasia officinalis</i> subsp. <i>monticola</i> | Montane Eyebright | Mountain pasture |
| <i>Euphrasia officinalis</i> subsp. <i>pratensis</i> | Rostkov's Eyebright | Valley meadows |
| <i>Euphrasia salisburgensis</i> | Irish Eyebright | Pasture |
| <i>Melampyrum pratense</i> | Common Cow-wheat | Woods |
| <i>Melampyrum sylvaticum</i> | Small Cow-wheat | Forest |
| <i>Orobanche alba</i> | Thyme Broomrape | Occasional |
| <i>Orobanche caryophyllacea</i> | Clove-scented Broomrape | Occasional |
| <i>Pedicularis foliosa</i> | Leafy Lousewort | Schynige Platte |
| <i>Pedicularis kernerii</i> | Kerner's Lousewort | Alpine turf |
| <i>Pedicularis oederi</i> | Crimson-tipped Lousewort | Alpine turf |
| <i>Pedicularis tuberosa</i> | Long-beaked Yellow Lousewort | Acidic turf |
| <i>Pedicularis verticillata</i> | Whorled Lousewort | Mountain turf |
| <i>Rhinanthus alectorolophus</i> | Greater Yellow-rattle | Common. Meadows |
| <i>Rhinanthus minor</i> | Yellow-rattle | Frequent in pasture |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---------------------------------|-----------------------------|----------------------------|
| Oxalidaceae | | |
| <i>Oxalis acetosella</i> | Wood-sorrel | Shaded rocks |
| Papaveraceae | | |
| <i>Chelidonium majus</i> | Greater Celadine | Occasional |
| <i>Papaver cambricum</i> ‡ | Welsh Poppy | Occasional |
| <i>Papaver rhoeas</i> | Common Poppy | Occasional. Lowland |
| Plantaginaceae | | |
| <i>Cymbalaria muralis</i> ‡ | Ivy-leaved Toadflax | Occasional on walls |
| <i>Erinus alpinus</i> | Fairy Foxglove | Frequent |
| <i>Globularia cordifolia</i> | Matted Globularia | Frequent |
| <i>Globularia nudicaulis</i> | Leafless-stemmed Globularia | Frequent |
| <i>Linaria alpina</i> | Alpine Toadflax | Moraine and scree |
| <i>Plantago alpina</i> | Alpine Plantain | Mountain turf |
| <i>Plantago atrata</i> | Dark Plantain | Mountain turf |
| <i>Plantago major</i> | Greater Plantain | Occasional |
| <i>Plantago serpentina</i> | Fleshy Plantain | Frequent |
| <i>Plantago media</i> | Hoary Plantain | Occasional |
| <i>Plantago lanceolata</i> | Ribwort Plantain | Occasional |
| <i>Veronica alpina</i> | Alpine Speedwell | Stony mountain pasture |
| <i>Veronica aphylla</i> | Leafless-stemmed Speedwell | Stony mountain pasture |
| <i>Veronica beccabunga</i> | Brooklime | Wet places |
| <i>Veronica bellidioides</i> | Violet Speedwell | Acidic pasture |
| <i>Veronica chamaedrys</i> | Germander Speedwell | Below Wengen |
| <i>Veronica fruticans</i> | Rock Speedwell | Mountain rocks |
| <i>Veronica fruticulosa</i> | Shrubby Speedwell | Oberberghorn |
| <i>Veronica officinalis</i> | Heath Speedwell | Woods |
| <i>Veronica serpyllifolia</i> | Thyme-leaved Speedwell | Frequent |
| <i>Veronica urticifolia</i> | Nettle-leaved Speedwell | Woods |
| Polygalaceae | | |
| <i>Polygala alpestris</i> | Mountain Milkwort | Common |
| <i>Polygala alpina</i> | Alpine Milkwort | Mountain rocks |
| <i>Polygala vulgaris</i> | Common Milkwort | Frequent |
| <i>Polygaloides chamaebuxus</i> | Shrubby Milkwort | Frequent on rocks |
| Polygonaceae | | |
| <i>Bistorta officinalis</i> | Common Bistort | Damp pasture |
| <i>Bistorta vivipara</i> | Alpine Bistort | Common. Mountain turf |
| <i>Oxyria digyna</i> | Mountain Sorrel | Mountain rocks and scree |
| <i>Reynoutria japonica</i> ‡ | Japanese Knotweed | Lowland |
| <i>Rumex acetosa</i> | Common Sorrel | Widespread |
| <i>Rumex acetosella</i> | Sheep's Sorrel | Acidic pasture |
| <i>Rumex alpestris</i> | Mountain Dock | Mountain pasture |
| <i>Rumex alpinus</i> | Monk's-rhubarb | Disturbed mountain pasture |
| <i>Rumex obtusifolius</i> | Broad-leaved Dock | Occasional |
| Plantain Family | | |
| Milkwort Family | | |
| Dock Family | | |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|----------------------------|-------------------------------|
| <i>Rumex scutatus</i> | French Sorrel | Rocks and scree |
| Primulaceae | | |
| <i>Androsace chamaejasme</i> | Ciliate Rock-jasmine | Frequent |
| <i>Androsace helvetica</i> | Swiss Rock-jasmine | Alpine cliffs |
| <i>Androsace obtusifolia</i> | Blunt-leaved Rock-jasmine | Acidic turf |
| <i>Androsace pubescens</i> | Downy Rock-jasmine | Rare. Mountain rocks |
| <i>Lysimachia nemorum</i> | Yellow Pimpernel | Damp woods |
| <i>Primula auricula</i> | Auricula | Mountain rocks and turf |
| <i>Primula elatior</i> | Oxlip | Snowbeds |
| <i>Primula farinosa</i> | Bird's-eye Primrose | Mountain turf |
| <i>Primula hirsuta</i> | Hairy Primrose | Acidic rocks |
| <i>Primula veris</i> | Cowslip | Below Wengen |
| <i>Soldanella alpina</i> | Alpine Snowbell | Common |
| Ranunculaceae | | |
| <i>Aconitum lycoctonum</i> subsp. <i>vulparia</i> | Wolf's-bane | Frequent. Forest |
| <i>Aconitum napellus</i> subsp. <i>napellus</i> | Compact Monk's-hood | Damp meadows |
| <i>Actaea spicata</i> | Baneberry | Woods |
| <i>Anemonastrum narcissiflorum</i> | Narcissus-flowered Anemone | Mountain pasture |
| <i>Anemonoides nemorosa</i> | Wood Anemone | Woods |
| <i>Aquilegia alpina</i> | Alpine Columbine | Rare. Biglenalp |
| <i>Aquilegia atrata</i> | Dark Columbine | Forest and meadows |
| <i>Caltha palustris</i> | Marsh-marigold | Flushes and streamsides |
| <i>Clematis vitalba</i> | Old Man's Beard | Common. Lowland |
| <i>Hepatica nobilis</i> | Liverleaf | Woodland |
| <i>Pulsatilla alpina</i> subsp. <i>alpina</i> | Alpine Pasqueflower | Frequent on limestone |
| <i>Pulsatilla alpina</i> subsp. <i>apiifolia</i> | Alpine Pasqueflower | Frequent on acidic soils |
| <i>Pulsatilla vernalis</i> | Spring Pasqueflower | Locally common |
| <i>Ranunculus aconitifolius</i> | Aconite-leaved Buttercup | Locally common |
| <i>Ranunculus acris</i> | Meadow Buttercup | Common |
| <i>Ranunculus alpestris</i> | Alpine Buttercup | Snowbeds. Common |
| <i>Ranunculus glacialis</i> | Glacier Crowfoot | Alpine screes. Eiger |
| <i>Ranunculus kuepferi</i> | Kupfer's Buttercup | Locally common |
| <i>Ranunculus montanus</i> | Mountain Buttercup | Mountain pasture |
| <i>Ranunculus repens</i> | Creeping Buttercup | Occasional. Meadows |
| <i>Thalictrum aquilegifolium</i> | Great Meadow-rue | Mountain meadows |
| <i>Thalictrum foetidum</i> | Stinking Meadow-rue | Trummelbach |
| <i>Trollius europaeus</i> | Globeflower | Common |
| Rhamnaceae | | |
| <i>Rhamnus alpinus</i> | Alpine Buckthorn | Rocky places |
| <i>Rhamnus pumilus</i> | Dwarf Buckthorn | Limestone rocks. Oberberghorn |
| Rosaceae | | |
| <i>Agrimonia eupatoria</i> | Agrimony | Occasional. Lowland |
| <i>Alchemilla alpina</i> | Alpine Lady's-mantle | Common |
| Buttercup Family | | |
| Rose Family | | |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|--|---------------------------|------------------------------|
| <i>Alchemilla conjuncta</i> | Silver Lady's-mantle | Frequent |
| <i>Alchemilla glabra</i> | Smooth Lady's-mantle | Frequent |
| <i>Alchemilla pentaphyllea</i> | Cut-leaved Lady's-mantle | Snowbeds |
| <i>Alchemilla vulgaris</i> agg. | Lady's-mantle | Frequent |
| <i>Alchemilla xanthochlora</i> | Pale Lady's-mantle | Frequent |
| <i>Aria edulis</i> | Whitebeam | Occasional. Woods |
| <i>Aruncus dioicus</i> | Goat's-beard Spiraea | Woods. Below Wengen |
| <i>Chamaemespilus alpina</i> | False-medlar | Mountain pasture |
| <i>Cotoneaster integerrimus</i> | Wild Cotoneaster | Mountain rocks. Oberberghorn |
| <i>Crataegus monogyna</i> | Hawthorn | Woods |
| <i>Dryas octopetala</i> | Mountain Avens | Mountain rocks and scree |
| <i>Filipendula ulmaria</i> | Meadowsweet | Wet meadows |
| <i>Fragaria vesca</i> | Wild Strawberry | Woods and pasture |
| <i>Geum montanum</i> | Alpine Avens | Mountain turf |
| <i>Geum rivale</i> | Water Avens | Occasional. Wet pasture |
| <i>Geum urbanum</i> | Herb Bennet | Woods |
| <i>Hedlundia mougeotii</i> | Mougeot's Whitebeam | Woods |
| <i>Potentilla aurea</i> | Golden Cinquefoil | Occasional. Mountain turf |
| <i>Potentilla crantzii</i> | Alpine Cinquefoil | Common. Mountain turf |
| <i>Potentilla erecta</i> | Tormentil | Acidic turf |
| <i>Potentilla grandiflora</i> | Large-flowered Cinquefoil | Mannlichen |
| <i>Potentilla sterilis</i> | Barren Strawberry | Frequent |
| <i>Poterium sanguisorba</i> | Salad Burnet | Widespread |
| <i>Prunus avium</i> | Wild Cherry | Woods |
| <i>Rosa canina</i> | Dog-rose | Below Wengen |
| <i>Rosa glauca</i> | Blue-leaved Rose | Occasional |
| <i>Rosa pendulina</i> | Alpine Rose | Frequent. Mountains |
| <i>Rosa rubiginosa</i> | Sweet-briar | Below Wengen |
| <i>Rubus fruticosus</i> agg. | Bramble | Below Wengen |
| <i>Rubus idaeus</i> | Raspberry | Woods |
| <i>Rubus saxatilis</i> | Stone Bramble | Occasional. Rocky places |
| <i>Sibbaldia procumbens</i> | Sibbaldia | Snowbeds |
| <i>Sorbus aucuparia</i> | Rowan | Woods |
| Rubiaceae | Madder Family | |
| <i>Cruciata laevipes</i> | Crosswort | Meadows |
| <i>Cynanchica pyrenaica</i> subsp. <i>cynanchica</i> | Squinancywort | Occasional |
| <i>Galium album</i> | White Bedstraw | Meadows |
| <i>Galium anisophyllum</i> | Alpine Bedstraw | Frequent |
| <i>Galium megalospermum</i> | Swiss Bedstraw | Moraines |
| <i>Galium odoratum</i> | Woodruff | Woods |
| <i>Galium pumilum</i> | Slender Bedstraw | Frequent |
| <i>Galium rotundifolium</i> | Round-leaved Bedstraw | Grutschalp |
| <i>Galium verum</i> | Lady's Bedstraw | Meadows |
| Salicaceae | Willow Family | |
| <i>Populus alba</i> | White Poplar | Lowland |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|--|---------------------------|
| <i>Populus tremula</i> | Aspen | Occasional |
| <i>Salix aurita</i> | Eared Willow | Widespread |
| <i>Salix breviserrata</i> | Finely-toothed Willow | Occasional |
| <i>Salix caprea</i> | Goat Willow | Lowland |
| <i>Salix eleagnos</i> | Hoary Willow | Lowland. By rivers |
| <i>Salix foetida</i> | Stinking Willow | Locally common |
| <i>Salix hastata</i> | Large-stipuled Willow | Local. Lauberhorn |
| <i>Salix glaucosericea</i> | Silky Willow | Local. Lauberhorn |
| <i>Salix herbacea</i> | Least Willow | Mountain turf |
| <i>Salix laggeri</i> | Lagger's Willow | Local. Lauberhorn |
| <i>Salix purpurea</i> | Purple Willow | Lowland |
| <i>Salix reticulata</i> | Net-leaved Willow | Alpine rocks and turf |
| <i>Salix retusa</i> | Retuse-leaved Willow | Alpine rocks and turf |
| <i>Salix serpillifolia</i> | Thyme-leaved Willow | Alpine rocks and turf |
| <i>Salix viminalis</i> | Osier | Lowland. Riversides |
| Santalaceae | | |
| <i>Thesium alpinum</i> | Bastard-toadflax Family Alpine Bastard-toadflax | Occasional. Pasture |
| Sapindaceae | | |
| <i>Acer campestre</i> | Maple Family Field Maple | Frequent |
| <i>Acer platanoides</i> | Norway Maple | Occasional |
| <i>Acer pseudoplatanus</i> | Sycamore | Common |
| Saxifragaceae | | |
| <i>Chrysosplenium alternifolium</i> | Saxifrage Family Alternate-leaved Golden-saxifrage | Occasional |
| <i>Micranthes stellaris</i> subsp. <i>robusta</i> | Starry Saxifrage | Acidic flushes |
| <i>Saxifraga aizoides</i> | Yellow Mountain Saxifrage | Wet turf and scree |
| <i>Saxifraga androsacea</i> | Scree Saxifrage | Damp scree and snowbeds |
| <i>Saxifraga aspera</i> | Rough Saxifrage | Occasional. Rocks |
| <i>Saxifraga bryoides</i> | Mossy Saxifrage | Acidic rocks |
| <i>Saxifraga caesia</i> | Blue Saxifrage | Limestone rocks and scree |
| <i>Saxifraga moschata</i> | Musky Saxifrage | Mountain rocks |
| <i>Saxifraga oppositifolia</i> | Purple Saxifrage | Rocks and moraine |
| <i>Saxifraga paniculata</i> | Livelong Saxifrage | Rocks |
| <i>Saxifraga rotundifolia</i> | Round-leaved Saxifrage | Damp shaded rocks |
| Scrophulariaceae | | |
| <i>Verbascum nigrum</i> | Figwort Family Dark Mullein | Lowland |
| Solanaceae | | |
| <i>Atropa bella-donna</i> | Nightshade Family Deadly Nightshade | Forest. Wilderswil |
| Thymelaeaceae | | |
| <i>Daphne mezereum</i> | Mezereon Family Mezereon | Local. Mountain pasture |
| Ulmaceae | | |
| | Elm Family | |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|--|------------------------------|--|
| <i>Ulmus glabra</i> | Wych Elm | Valley woods |
| Urticaceae | Nettle Family | |
| <i>Urtica dioica</i> | Common Nettle | Widespread |
| Viburnaceae | Viburnum Family | |
| <i>Adoxa moschatellina</i> | Moschatel | Rare. Mannlichen |
| <i>Sambucus ebulus</i> | Dwarf Elder | Lowland. Wilderswil |
| <i>Sambucus nigra</i> | Common Elder | Woods |
| <i>Sambucus racemosa</i> | Alpine Elder | Occasional. Rocky places |
| <i>Viburnum lantana</i> | Wayfaring-tree | Woods below Wengen |
| <i>Viburnum opulus</i> | Guelder-rose | Woods below Wengen |
| Violaceae | Violet Family | |
| <i>Viola biflora</i> | Yellow Wood Violet | Common |
| <i>Viola calcarata</i> | Long-spurred Pansy | Locally common |
| <i>Viola cenisia</i> | Mt Cenis Pansy | Moraine and scree |
| <i>Viola lutea</i> | Mountain Pansy | Locally common |
| <i>Viola palustris</i> | Marsh Violet | Common |
| <i>Viola riviniana</i> | Common Dog-violet | Common |
| <i>Viola tricolor</i> subsp. <i>tricolor</i> | Wild Pansy | Local |
| MONOCOTYLEDONS | | |
| Amaryllidaceae | Daffodil Family | |
| <i>Allium schoenoprasum</i> | Chives | Occasional. Mountains Uncommon. Meadows. Trummelbach |
| <i>Allium scordoprasum</i> | Sand Leek | Mannlichen |
| <i>Allium lusitanicum</i> | Mountain Onion | Grutschalp, Schynige Platte |
| <i>Allium victorialis</i> | Alpine Leek | |
| Asparagaceae | Asparagus Family | |
| <i>Anthericum liliago</i> | St Bernard's Lily | Meadow above Wengen |
| <i>Anthericum ramosum</i> | Branched St Bernard's Lily | Trummelbach |
| <i>Convallaria majalis</i> | Lily-of-the-Valley | Limestone cliffs and scree |
| <i>Maianthemum bifolium</i> | May Lily | Mountain woods |
| <i>Paradisea liliastrum</i> | St Bruno's Lily | Local. Mountain pasture |
| <i>Polygonatum odoratum</i> | Angular Solomon's-seal | Limestone cliffs and scree |
| <i>Polygonatum verticillatum</i> | Whorled Solomon's-seal | Occasional. Forests |
| Colchicaceae | Meadow Saffron Family | |
| <i>Colchicum autumnale</i> | Meadow Saffron | Meadows below Wengen |
| Cyperaceae | Sedge Family | |
| <i>Carex atrata</i> subsp. <i>atrata</i> | Black Alpine Sedge | Wet mountain pasture |
| <i>Carex canescens</i> | White Sedge | Wet mountain pasture |
| <i>Carex capillaris</i> | Hair Sedge | Wet mountain pasture |
| <i>Carex caryophylla</i> | Spring Sedge | Dry pasture |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---------------------------------|---|--------------------------------|
| <i>Carex curvula</i> | An alpine Sedge | Dominant. Mountain turf |
| <i>Carex davalliana</i> | Davall's Sedge | Wet calcareous turf |
| <i>Carex dioica</i> | Dioecious Sedge | Base-rich flushes |
| <i>Carex echinata</i> | Star Sedge | Frequent |
| <i>Carex ferruginea</i> | Rusty Sedge | Dominant. Limestone turf |
| <i>Carex flacca</i> | Glaucous Sedge | Common |
| <i>Carex flava</i> agg. | Yellow-sedge | Marshes |
| <i>Carex nigra</i> | Common Sedge | Common |
| <i>Carex panicea</i> | Carnation Sedge | Common |
| <i>Carex rostrata</i> | Bottle Sedge | Pond margins |
| <i>Carex sempervirens</i> | Evergreen Sedge | Locally dominant on poor soils |
| <i>Eriophorum angustifolium</i> | Common Cottongrass | Wet flushes and bogs |
| <i>Eriophorum latifolium</i> | Broad-leaved Cottongrass | Marshes |
| <i>Eriophorum scheuchzeri</i> | Scheuchzer's Cottongrass | Occasional. Bogs |
| <i>Eriophorum vaginatum</i> | Hare's-tail Cottongrass | Acidic bogs |
| <i>Trichophorum cespitosum</i> | Northern Deergrass | Acidic bogs. Uncommon |
| Iridaceae | | |
| <i>Crocus vernus</i> | Iris Family Spring Crocus | Common |
| Juncaceae | | |
| <i>Juncus alpinoarticulatus</i> | Rush Family Alpine Rush | Wet mountain pasture |
| <i>Juncus effusus</i> | Soft Rush | Wet pasture |
| <i>Juncus inflexus</i> | Hard Rush | Wet pasture |
| <i>Juncus jacquinii</i> | Jacquin's Rush | Wet mountain pasture |
| <i>Juncus triglumis</i> | Three-flowered Rush | Acidic mountain turf |
| <i>Luzula alpinopilosa</i> | Alpine Wood-rush | Mountain turf |
| <i>Luzula lutea</i> | Yellow Wood-rush | Mountain turf |
| <i>Luzula multiflora</i> | Heath Wood-rush | Mountain turf |
| <i>Luzula spicata</i> | Spiked Wood-rush | Mountain turf |
| <i>Luzula sylvatica</i> | Great Wood-rush | Woods |
| <i>Oreojuncus trifidus</i> | Three-leaved Rush | Acidic mountain turf |
| Liliaceae | | |
| <i>Gagea fragifera</i> | Lily Family A Yellow Star-of-Bethlehem | Uncommon. Mountain turf |
| <i>Gagea serotina</i> | Snowdon Lily | Mountain turf |
| <i>Lilium martagon</i> | Martagon Lily | Woods |
| <i>Streptopus amplexifolius</i> | Streptopus | Occasional. Forests |
| Melanthiaceae | | |
| <i>Paris quadrifolia</i> | Herb-Paris Family Herb-Paris | Local. Woods |
| <i>Veratrum album</i> | White False-helleborine | Mountain pasture |
| Orchidaceae | | |
| <i>Cephalanthera longifolia</i> | Orchid Family Narrow-leaved Helleborine | Frequent. Woods |
| <i>Chamorchis alpina</i> | Dwarf Orchid | Local |
| <i>Coeloglossum viride</i> | Frog Orchid | Frequent |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|---|---------------------------|----------------------------|
| <i>Corallorhiza trifida</i> | Coralroot Orchid | Local. Forests |
| <i>Cypripedium calceolus</i> | Lady's-slipper | Local |
| <i>Dactylorhiza maculata</i> subsp. <i>fuchsii</i> | Common Spotted-orchid | Damp base-rich soils |
| <i>Dactylorhiza maculata</i> subsp. <i>maculata</i> | Heath Spotted-orchid | Acidic soils |
| <i>Dactylorhiza majalis</i> | Broad-leaved Marsh-orchid | Common |
| <i>Epipactis atrorubens</i> | Dark-red Helleborine | Frequent. Forests |
| <i>Epipactis helleborine</i> | Broad-leaved Helleborine | Frequent. Forests |
| <i>Gymnadenia conopsea</i> | Fragrant-orchid | Common |
| <i>Neotinea ustulata</i> | Burnt Orchid | Uncommon. Mountain pasture |
| <i>Neottia cordata</i> | Lesser Twayblade | Locally common |
| <i>Neottia nidus-avis</i> | Bird's-nest Orchid | Frequent. Forests |
| <i>Neottia ovata</i> | Twayblade | Locally common |
| <i>Nigritella rhellicani</i> | Alpine Vanilla Orchid | Locally common |
| <i>Orchis mascula</i> | Early-purple Orchid | Common |
| <i>Platanthera bifolia</i> | Lesser Butterfly-orchid | Locally frequent. Forests |
| <i>Platanthera chlorantha</i> | Greater Butterfly-orchid | Local. Woods |
| <i>Pseudorchis albida</i> | Small-white Orchid | Common. Mountain turf |
| <i>Traunsteinera globosa</i> | Round-headed Orchid | Local |

Poaceae

| Latin Name | Common Name | Notes |
|--------------------------------|---------------------|----------------------------|
| <i>Agrostis capillaris</i> | Common Bent | Common |
| <i>Anthoxanthum alpinum</i> | Alpine Vernal-grass | Frequent. Mountain turf |
| <i>Anthoxanthum odoratum</i> | Sweet Vernal-grass | Common |
| <i>Arrhenatherum elatius</i> | False Oat-grass | Common |
| <i>Avenella flexuosa</i> | Wavy Hair-grass | Acidic mountain turf |
| <i>Brachypodium sylvaticum</i> | False Brome | Woods |
| <i>Briza media</i> | Quaking-grass | Locally common |
| <i>Bromus hordeaceus</i> | Soft Brome | Frequent |
| <i>Bromus sterilis</i> | Barren Brome | Frequent |
| <i>Cynosurus cristatus</i> | Crested Dog's-tail | Common |
| <i>Dactylis glomerata</i> | Cock's-foot | Common |
| <i>Deschampsia cespitosa</i> | Tufted Hair-grass | Wet mountain pasture |
| <i>Festuca ovina</i> | Sheep's-fescue | Common |
| <i>Festuca rubra</i> | Red Fescue | Common |
| <i>Festuca violacea</i> | Violet Fescue | Mountain turf |
| <i>Holcus lanatus</i> | Yorkshire-fog | Locally common |
| <i>Holcus mollis</i> | Creeping Soft-grass | Locally common |
| <i>Koeleria macrantha</i> | Crested Hair-grass | Frequent |
| <i>Lolium perenne</i> | Perennial Rye-grass | Frequent |
| <i>Melica nutans</i> | Mountain Melick | Woods |
| <i>Nardus stricta</i> | Mat-grass | Common |
| <i>Phleum alpinum</i> | Alpine Timothy | Frequent. Mountain pasture |
| <i>Phleum pratense</i> | Timothy | Common |
| <i>Poa alpina</i> | Alpine Meadow-grass | Common |
| <i>Poa annua</i> | Annual Meadow-grass | Common on trackways |
| <i>Poa pratensis</i> | Smooth Meadow-grass | Common. Meadows |
| <i>Poa nemoralis</i> | Wood Meadow-grass | Frequent |

‡ = cultivated or naturalised

| Latin Name | Common Name | Notes |
|------------------------------|--------------------------|---------------------------|
| <i>Poa supina</i> | Creeping Meadow-grass | Frequent |
| <i>Poa trivialis</i> | Rough Meadow-grass | Frequent |
| <i>Schedonorus pratensis</i> | Meadow Fescue | Common |
| <i>Sesleria caerulea</i> | Blue Moor-grass | Common. Limestone pasture |
| Tofieldiaceae | | |
| <i>Tofieldia calyculata</i> | Scottish Asphodel Family | Common |

Birds (H = Heard only)

| Common name | Latin name | June/July 2025 | | | | | | | |
|--------------------------|--------------------------------|----------------|----|----|----|----|----|---|---|
| | | 25 | 26 | 27 | 28 | 29 | 30 | 1 | 2 |
| Common Swift | <i>Apus apus</i> | ✓ | ✓ | ✓ | | ✓ | ✓ | ✓ | |
| Common Cuckoo | <i>Cuculus canorus</i> | | | | H | | | | |
| Common Wood Pigeon | <i>Columba palumbus</i> | | ✓ | | | | | | |
| Common Buzzard | <i>Buteo buteo</i> | | | | | | ✓ | | |
| Tawny Owl | <i>Strix aluco</i> | | | H | | H | | | |
| Great Spotted Woodpecker | <i>Dendrocopos major</i> | | | | | | | H | |
| Eurasian Jay | <i>Garrulus glandarius</i> | ✓ | | | | | | | |
| Eurasian Magpie | <i>Pica pica</i> | | H | ✓ | ✓ | ✓ | ✓ | ✓ | |
| Northern Nutcracker | <i>Nucifraga caryocatactes</i> | | | ✓ | ✓ | ✓ | ✓ | H | |
| Alpine Chough | <i>Pyrrhocorax graculus</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| Carrion Crow | <i>Corvus corone</i> | | ✓ | | | | | ✓ | |
| Northern Raven | <i>Corvus corax</i> | | | ✓ | | ✓ | | | |
| Coal Tit | <i>Parus ater</i> | | | | | ✓ | | | |
| Eurasian Blue Tit | <i>Cyanistes caeruleus</i> | | ✓ | | | | ✓ | | |
| Great Tit | <i>Parus major</i> | | ✓ | ✓ | ✓ | | ✓ | ✓ | |
| Eurasian Skylark | <i>Alauda arvensis</i> | | | | H | H | | | |
| Barn Swallow | <i>Hirundo rustica</i> | | ✓ | | | | | | |
| Western House Martin | <i>Delichon urbicum</i> | | ✓ | | | | | | |
| Common Chiffchaff | <i>Phylloscopus collybita</i> | | | | | H | | | |
| Eurasian Blackcap | <i>Sylvia atricapilla</i> | | ✓ | ✓ | | H | | | |
| Garden Warbler | <i>Sylvia borin</i> | | | ✓ | | | | | |
| Lesser Whitethroat | <i>Curruca curruca</i> | | | | | ✓ | H | | |
| Eurasian Wren | <i>Troglodytes troglodytes</i> | | ✓ | | | | ✓ | | |
| Common Starling | <i>Sturnus vulgaris</i> | | | | | | | ✓ | |
| Song Thrush | <i>Turdus philomelos</i> | | H | ✓ | | ✓ | | | |
| Common Blackbird | <i>Turdus merula</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Fieldfare | <i>Turdus pilaris</i> | | | | | | H | | |
| Ring Ouzel | <i>Turdus torquatus</i> | | | | H | | ✓ | | |
| Black Redstart | <i>Phoenicurus ochruros</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| White-winged Snowfinch | <i>Montifringilla nivalis</i> | | | | ✓ | | | | |
| House Sparrow | <i>Passer domesticus</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Dunnock | <i>Prunella modularis</i> | | | | | | ✓ | | |
| Grey Wagtail | <i>Motacilla cinerea</i> | | ✓ | | | H | | | |
| White Wagtail | <i>Motacilla alba</i> | | | ✓ | ✓ | ✓ | | | |
| Water Pipit | <i>Anthus spinoletta</i> | | | | ✓ | ✓ | ✓ | ✓ | |
| Eurasian Chaffinch | <i>Fringilla coelebs</i> | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ |
| Common Linnet | <i>Linaria cannabina</i> | | ✓ | | | | | | |

| | | June/July 2025 | | | | | | | |
|--------------------|----------------------------|----------------|----|----|----|----|----|---|---|
| Common name | Latin name | 25 | 26 | 27 | 28 | 29 | 30 | 1 | 2 |
| European Goldfinch | <i>Carduelis carduelis</i> | | ✓ | ✓ | | | | | |

Vertebrates

| | | June/July 2025 | | | | | | | |
|-----------------------|----------------------------|----------------|----|----|----|----|----|---|---|
| Common name | Latin name | 25 | 26 | 27 | 28 | 29 | 30 | 1 | 2 |
| Mammals | | | | | | | | | |
| Eurasian Red Squirrel | <i>Sciurus vulgaris</i> | | ✓ | | | | | | |
| Roe Deer | <i>Capreolus capreolus</i> | ✓ | | | | | ✓ | | |
| Alpine Chamois | <i>Rupicapra rupicapra</i> | | | | 24 | | | | |
| Amphibians | | | | | | | | | |
| Common Frog | <i>Rana temporaria</i> | | | ✓ | | ✓ | | | |
| Common Newt | <i>Triturus vulgaris</i> | | | ✓ | ✓ | | | | |

Invertebrates

| | | June/July 2025 | | | | | | | |
|---------------------------------|---------------------------------|----------------|----|----|----|----|----|---|---|
| Common name | Latin name | 25 | 26 | 27 | 28 | 29 | 30 | 1 | 2 |
| Butterflies | | | | | | | | | |
| Papilionidae | | | | | | | | | |
| Swallowtail | <i>Papilio machaon</i> | | | ✓ | ✓ | ✓ | ✓ | ✓ | |
| Hesperiidae | | | | | | | | | |
| Essex Skipper | <i>Thymelicus lineola</i> | | ✓ | | | | | | |
| Alpine Grizzled Skipper | <i>Pyrgus andromedae</i> | | | | ✓ | ✓ | | | |
| Chequered Skipper | <i>Carterocephalus palaemon</i> | | | ✓ | | | | | |
| Pieridae | | | | | | | | | |
| Clouded Yellow | <i>Colias crocea</i> | | | | | ✓ | ✓ | | |
| Black-veined White | <i>Aporia crataegi</i> | | | ✓ | | | | | |
| Peak White | <i>Pontia callidice</i> | | | | | | | ✓ | |
| Large White | <i>Pieris brassicae</i> | | ✓ | ✓ | ✓ | | ✓ | ✓ | |
| Small White | <i>Pieris rapae</i> | | ✓ | ✓ | ✓ | | | ✓ | |
| Orange Tip | <i>Anthocharis cardamines</i> | | | ✓ | | ✓ | | | |
| Lycaenidae | | | | | | | | | |
| Little Blue | <i>Cupido minimus</i> | | | ✓ | ✓ | ✓ | | | |
| Mazarine Blue | <i>Cyaniris semiargus</i> | | | ✓ | ✓ | ✓ | | ✓ | |
| Common Blue | <i>Polyommatus icarus</i> | | | | | ✓ | ✓ | | |
| Alpine Blue | <i>Agriades orbitulus</i> | | | ✓ | | | | | |
| Nymphalidae | | | | | | | | | |
| Mountain Fritillary | <i>Boloria napaea</i> | | | | | ✓ | | | |
| Small Pearl-bordered Fritillary | <i>Boloria selene</i> | | | ✓ | | ✓ | | ✓ | |
| Pearl-bordered Fritillary | <i>Boloria euphrosyne</i> | | | ✓ | | | | | |
| Painted Lady | <i>Vanessa cardui</i> | | | ✓ | ✓ | ✓ | | ✓ | |
| Red Admiral | <i>Vanessa atalanta</i> | | ✓ | ✓ | | ✓ | | ✓ | |
| Small Tortoiseshell | <i>Aglais urticae</i> | | | ✓ | ✓ | | ✓ | ✓ | |
| False Heath Fritillary | <i>Melitaea diamina</i> | | | | | ✓ | | | |
| Alpine Heath | <i>Coenonympha gardetta</i> | | | ✓ | | ✓ | | | |

| Common name | Latin name | June/July 2025 | | | | | | | |
|--------------------------|---------------------------------|----------------|----|----|----|----|----|---|---|
| | | 25 | 26 | 27 | 28 | 29 | 30 | 1 | 2 |
| Marbled White | <i>Melanargia galathea</i> | | ✓ | ✓ | | ✓ | | | |
| Meadow Brown | <i>Maniola jurtina</i> | | ✓ | | | | | | |
| Piedmont Ringlet | <i>Erebia meolans</i> | | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| | | | | | | | | | |
| Moths | | | | | | | | | |
| Oak Eggar | <i>Lasiocampa quercus</i> | | | | | | ✓ | | |
| Latticed Heath | <i>Chiasmia clathrata</i> | | | | | | ✓ | | |
| a Geometrid | <i>Psodos quadrifaria</i> | | | | ✓ | | ✓ | | |
| White-spotted Sable | <i>Anania funebris</i> | | | ✓ | | | | | |
| Common Heath | <i>Ematurga atomaria</i> | | | ✓ | ✓ | | | | |
| Silver-ground Carpet | <i>Xanthorhoe montanata</i> | | | | ✓ | | | | |
| | | | | | | | | | |
| Dragonflies | | | | | | | | | |
| Common Blue Damselfly | <i>Enallagma cyathigerum</i> | | | | | ✓ | | | |
| Broad-bodied Chaser | <i>Libellula depressa</i> | | | ✓ | | | | | |
| Four-spotted Chaser | <i>Libellula quadrimaculata</i> | | | ✓ | | | | | |
| White-faced Darter | <i>Leucorrhinia dubia</i> | | | | | ✓ | | | |
| Blue Emperor | <i>Anax imperator</i> | | | | ✓ | | | | |
| Southern Hawker | <i>Aeshna cyanea</i> | | | | | ✓ | | | |
| | | | | | | | | | |
| Other Insects | | | | | | | | | |
| Great Green Bush-cricket | <i>Tettigonia viridissima</i> | | | | | ✓ | | | |
| Red Wood Ant | <i>Formica rufa</i> | | | ✓ | | ✓ | | | |
| Violet Oil Beetle | <i>Meloe violaceus</i> | | | | ✓ | | | | |